2020-2021 Synagis® Season – Prior Authorization Form

superior healthplan.

Today's Date: _____

Date Medication Required: _____

Phone: 1-800-218-7453 ext. 22080 | Fax: 1-866-683-5631

Section I — Dispensing Pharmacy Information

Name of Pharmacy	National Provider Identifier (NPI)	Area Code and Telephone N	o. Area Code and Fax No.	
Section II — Patient Demographics				
Name of Patient	Medicaid ID	Date of Birth (MMDDYY)	Gestational Age	
			weeks and / 7th day	
Address of Patient (Street, City, State, ZIP Coc	de) Patient Phone Number		County of Residence	
Has patient received a Synagis prophylactic injection during hospitalization since the start current of the RSV season?				
No Yes If yes, number of shots: Dose (mg): Date(s):				
Has the patient been hospitalization due to RS	V at any time since the start of the c	current RSV season?		
Section III — Patient Diagnosis at the start of the RSV season (Diagnosis/conditions must be clearly documented in the client's medical record.)				
Patients who are younger than 24 month chronological age can qualify, for up to five monthly doses of Synagis, based on diagnosis listed to the right.		ompromised during the RSV se nsplant, chemotherapy or other ompromised):		
	ICD-10-CM code:			
 Patients who are between 12 - 24 months chronological age at the start of the RSV season can qualify, for up to five monthly doses of Synagis, based on the diagnosis or conditions listed to the right. Please refer to page 2 for definition. 	of the following therapies wi that apply):	ithin the 6 months prior to the opsteroids $\square > 21\%$ Supplem	rematurity [#] , AND required any current RSV season (check all ental oxygen	
	24-3: Diagnosis of cystic fib or length less than the 10th	prosis with severe lung disease percentile:	*, or cystic fibrosis with weight	
	ICD-10-CM code:			
Patients who are younger than 12 months chronological age at the start of the RSV season can qualify, for up to five monthly doses of Synagis, based on criteria listed to the right.	□ 12-1: ≤ 28 6/7 weeks gestational age at birth:			
	ICD-10-CM code:	ICD-10-CM code:		
	12-2 : Chronic lung disease (CLD) of prematurity#:			
	ICD-10-CM code:			
	12-3 : Severe congenital abnormality of airway OR severe neuromuscular disease that impairs the ability to clear secretions from the upper airway because of ineffective cough:			
	ICD-10-CM code:			

2020-2021 Synagis® Season – Prior Authorization Form

	12-4 : Active diagnosis of hemodynamically significant congenital heart disease (CHD):	
	ICD-10-CM code: AND any of the below	
	Moderate to severe pulmonary hypertension.	
	Acyanotic heart disease, on medication to control congestive heart failure, and will require cardiac surgery.	
	Cyanotic heart disease.	
	(NOTE: This excludes infants with hemodynamically insignificant heart disease - refer to pages 2 and 3 for list.)	
	12-5: Diagnosis of cystic fibrosis with clinical evidence of CLD and/or nutritional compromise.	
	ICD-10-CM code:	
Section IV — Synagis Prescription detail (to be	completed by prescriber) Prescriber should send a prescription to the specialty pharmacy.	
Rx: Synagis (palivizumab) Injection Quan	tity: Dose (mg):	
Sig: Inject 15mg/kg one time per month Curre	nt Weight:(kg) or(lbs.) G 1*Epinephrine 1:1000 amp. Sig: Injected 0.01 mg/kg as directed.	
Prescriber Name	License No. NPI	
Address of Prescriber (Street, City, State and ZIP	Code) Area Code and Telephone No. Area Code and Fax No.	
Physician Signature	Date	

Fax the completed prior authorization form to Superior HealthPlan at 1-866-683-5631

Subcategories	
• Infants born < 32 weeks, 0 days' gestational age who require >21% oxygen for at least 28 days after birth.	
 Congestive heart failure (CHF) requiring medication Moderate to severe pulmonary hypertension Unrepaired cyanotic congenital heart disease 	
• Previous hospitalization for pulmonary exacerbation in the first year of life or abnormalities on chest radiography or chest computed tomography that persist when stable	
SED risk of RSV and generally should not receive immunoprophylaxis:	
 Secundum atrial septal defect Small ventriculoseptal defect Pulmonic stenosis Uncomplicated aortic stenosis Mild coarctation of the aorta Patent ductus arteriosus 	

3. Mild cardiomyopathy that does not require medical therapy for the condition.

2020-2021 Synagis® Season – Prior Authorization Form

Subcategories

4. Children in the second year of life on the basis of a history of prematurity alone.

Note: Tobacco smoke exposure is <u>not</u> an indication for Synagis administration. Tobacco dependent parents should be offered tobacco dependence treatment or referral for tobacco dependence treatment. 877-YES-QUIT (877-937-7848, YesQuit.org) is the Quitline operated in Texas.

Additional Information

- Texas Medicaid has adopted the updated guidance published in 2014 by the American Academy of Pediatrics.
- Infants born at 29 weeks, 0 days' gestation or later are no longer universally recommended to receive prophylaxis with Synagis.
- Infants born at 29 weeks, 0 days' gestation or later, on the basis of chronic lung disease, congenital heart disease, or another condition, may qualify to receive prophylaxis.
- Synagis is not recommended in the second year of life on the basis of prematurity alone.
- Monthly prophylaxis should be discontinued in any child who experiences a breakthrough RSV hospitalization.

References

Category

- "Updated Guidance for Palivizumab Prophylaxis Among Infants and Young Children at Increased Risk of Hospitalization for Respiratory Syncytial Virus Infection." *Pediatrics 134.*2 (2014): 415-420. Web. 11 Aug. 2015.
- Synagis® (palivizumab) [prescribing information]. Gaithersburg, MD: Medimmune, LLC. 2014.
- Epinephrine 1:1000 (1mg/ml) [prescribing information]. Lake Forest, IL: Hospira. 2008.