

Clinical Policy: Opioid Analgesics*

Reference Number: HIM.PA.139 Effective Date: 08.01.18 Last Review Date: 05.24 Line of Business: HIM

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

*Requests for transmucosal immediate-release fentanyl products (TIRFs) should be evaluated using the Fentanyl IR (Actiq, Fentora, Lazanda, Subsys) policy – CP.PMN.127.

Description

Opioid analgesics exert their analgesic effect through opiate receptors distributed in tissues throughout the body.

This policy applies to all formulary long and short acting opioids requiring prior authorization (PA) or any non-formulary opioid request.

FDA Approved Indication(s)

Opioid analgesics are indicated for the management and treatment of moderate to severe pain.

Policy/Criteria

Provider must submit documentation (including such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

I. Initial Approval Criteria

Please note: For HIM-Arkansas – if a member's covered prescription pain medication requires a prior authorization, then the prior authorization shall not be denied if the member has a terminal illness.

A. Cancer, Sickle Cell Disease, or Palliative Care (must meet all):

*Requests for transmucosal immediate-release fentanyl products (TIRFs) should be evaluated using the Fentanyl IR (Actiq, Fentora, Lazanda, Subsys) policy – CP.PMN.127.

- 1. Prescribed for pain associated with one of the following (a, b, or c):
 - a. Cancer;
 - b. Sickle cell disease;
 - c. Palliative care (hospice or any terminal condition);
- 2. Member has failed an adequate trial of two formulary short-acting opioid analgesics that does not require PA, dosed around the clock, unless clinically significant adverse effects are experienced, or all are contraindicated;

Approval duration: 12 months

- **B.** Short-Acting Agents Requests for \leq a 14-day Supply (must meet all):
 - 1. Prescribed for the treatment of pain unrelated to cancer, palliative care, or sickle cell disease;



- 2. Member has failed an adequate trial of two formulary short-acting opioids analgesics that does not require PA, dosed around the clock, unless clinically significant adverse effects are experienced, or all are contraindicated;
- For OHIO requests ONLY: Total opioid dose does not exceed 80 MME per day or for members who are stable (history of > 7 days of therapy) on doses higher than 80 MME per day, one of the following is met (a or b):
 - a. Provider will initiate a dose taper; *Future approval will require decrease from current dose.
 - b. Documentation that a dose taper has been attempted within the past 6 months, with the reasons for taper failure;
- 4. For OHIO requests ONLY: If opioid is being prescribed concomitantly with a benzodiazepine, member meets both of the following (a and b):
 - a. Documentation that the provider has acknowledged combined use of opioid medication and benzodiazepine;
 - b. Prescribed for short term (< 3 months) use or provider will discontinue concurrent use of benzodiazepine and opioid therapy within a 3-month period. **Re-authorization request for concurrent use of opioid and benzodiazepine will not be approved.*

Approval duration: 14 days

- C. Long-Acting Agents <u>OR</u> Requests Exceeding a 14-day Supply Within 28 Days <u>OR</u> Requests Exceeding a 28-day Supply Within 90 Days (must meet all):
 - 1. Prescribed for the treatment of pain unrelated to cancer, palliative care, or sickle cell disease;
 - 2. Member meets ALL of the following (a, b, and c), unless clinically significant adverse effects are experienced, or all are contraindicated:
 - a. Failure ≥ 2 non-opioid ancillary treatments (e.g., non-steroidal anti-inflammatory drugs [NSAIDs], acetaminophen, anticonvulsants, antidepressants);
 - b. For short-acting agent requests, one of the following (i or ii):
 - i. Prescribed agent is a formulary short-acting agent that does not require PA;
 - ii. Failure of an adequate trial of two formulary short-acting opioids analgesics, dosed around the clock;
 - c. For long-acting agent requests, <u>both</u> of the following (i and ii):
 - i. Except Louisiana ONLY, failure of an adequate trial of two short-acting opioids analgesics, dosed around the clock;
 - ii. Failure of an adequate trial of two formulary long-acting agents*; *For Louisiana, if request is for an abuse deterrent formulation, substitution shall not be made to an extended-release medication that does not have defined abuse deterrent properties.
 - For OHIO requests ONLY: Total opioid dose does not exceed 80 MME per day or for members who are stable (history of > 7 days of therapy) on doses higher than 80 MME per day, one of the following is met (a or b):
 - a. Provider will initiate a dose taper; *Future approval will require decrease from current dose.
 - b. Documentation that a dose taper has been attempted within the past 6 months, with the reasons for taper failure;



- 4. For OHIO requests ONLY: If opioid is being prescribed concomitantly with a benzodiazepine, member meets both of the following (a and b):
 - a. Documentation that the provider has acknowledged combined use of opioid medication and benzodiazepine;
 - b. Prescribed for short term (< 3 months) use or provider will discontinue concurrent use of benzodiazepine and opioid therapy within a 3-month period.

Approval duration:

Short-acting agents – 3 months or duration of request (whichever is less) **Long-acting agents** – 12 months

D. Diabetic Peripheral Neuropathy (must meet all):

- 1. Request is for Nucynta ER;
- 2. Diagnosis of diabetic peripheral neuropathy;
- 3. Age \geq 18 years;
- 4. Failure of gabapentin at ≥ 1,800 mg/day, unless contraindicated or clinically significant adverse effects are experienced;
- 5. Failure of a formulary tricyclic antidepressant (TCA) (e.g., amitriptyline, nortriptyline, imipramine) at up to maximally indicated doses, unless clinically significant adverse effects are experienced, or all are contraindicated;
- 6. Failure of a formulary serotonin-norepinephrine reuptake inhibitor (SNRI) (e.g., duloxetine, venlafaxine) at up to maximally indicated doses, unless clinically significant adverse effects are experienced, or all are contraindicated;
- 7. Dose does not exceed 500 mg per day.

Approval duration: 6 months or duration of request (whichever is less)

E. Other diagnoses/indications – Not applicable

II. Continued Therapy

Please note: For HIM-Arkansas – if a member's covered prescription pain medication requires a prior authorization, then the prior authorization shall not be denied if the member has a terminal illness.

A. Cancer, Sickle Cell Disease, Palliative Care (must meet all):

*Requests for transmucosal immediate-release fentanyl products (TIRFs) should be evaluated using the Fentanyl IR (Actiq, Fentora, Lazanda, Subsys) policy – CP.PMN.127.

 Currently receiving prescribed agent via Centene benefit for cancer, sickle cell disease, and palliative care or have previously met initial approval criteria;
Approval duration: 12 months





- **B.** Short-Acting Agents Requests for \leq a 14-day Supply (must meet all):
 - 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
 - 2. For OHIO requests ONLY: Total opioid dose does NOT exceed 80 MME/day or if the current dose is higher than 80 MME/day, one of the following is met (a, b or c):
 - a. Dose reduction has occurred since previous approval;
 - b. A dose taper has been attempted within the past 6 months and was not successful; **Reason(s) for taper failure must be provided*
 - c. Prescribed by or in consultation with a pain management specialist;
 - 3. For OHIO requests ONLY: If the requested dose is for > 80 MME/day, an increase in dose has not occurred since previous approval;
 - 4. For OHIO requests ONLY: If opioid is being prescribed concomitantly with a benzodiazepine, member meets all of the following (a, b, and c):
 - a. Currently receiving concurrent opioid and benzodiazepine therapy via Centene benefit or member has previously met the initial approval criteria;
 - b. Documentation supports that discontinuation of combination opioid and benzodiazepine therapy has been attempted in the last 3 months without success;
 - c. Prescribed by or in consultation with a pain management specialist.

Approval Duration: 14 days

- C. Long-Acting Agents <u>OR</u> Requests Exceeding a 14-day Supply Within 28 Days <u>OR</u> 28-day Supply Within 90 Days (must meet all):
 - 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (*refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B*);
 - 2. Has received more than a 14-day supply of opioid within 28 days or a 28-day supply within 90 days;

*If member does not meet this requirement, please use the initial approval criteria to review this request

- 3. Member continues to need opioid analgesics as evidenced by, including but not limited to <u>any</u> of the following:
 - a. Provider submits medical justification;
 - b. Documentation of recent (within the last 6 months) office visit or office chart notes demonstrating follow-up with the member;
 - c. Attestation that provider has reviewed the treatment plan with the member and assessed the risks and benefits of opioid dose and duration;
- 4. For OHIO requests ONLY: Total opioid dose does NOT exceed 80 MME/day or if the current dose is higher than 80 MME/day, one of the following is met (a, b, or c):
 - a. Dose reduction has occurred since previous approval;



- b. A dose taper has been attempted within the past 6 months and was not successful; **Reason(s) for taper failure must be provided*
- c. Prescribed by or in consultation with a pain management specialist;
- 5. For OHIO requests ONLY: If the requested dose is for > 80 MME/day, an increase in dose has not occurred since previous approval;
- 6. For OHIO requests ONLY: If opioid is being prescribed concomitantly with a benzodiazepine, member meets all of the following (a, b, and c):
 - a. Currently receiving concurrent opioid and benzodiazepine therapy via Centene benefit or member has previously met the initial approval criteria;
 - b. Documentation supports that discontinuation of combination opioid and benzodiazepine therapy has been attempted in the last 3 months without success;
 - c. Prescribed by or in consultation with a pain management specialist.

Approval duration:

Short-acting agents – 3 months or duration of request (whichever is less) **Long-acting agents** – 12 months

D. Diabetic Peripheral Neuropathy (must meet all):

- 1. Request is for Nucynta ER;
- 2. Currently receiving Nucynta ER for the diagnosis of diabetic peripheral neuropathy or member has met initial approval criteria;
- 3. Member continues to need Nucynta ER as evidenced by, including but not limited to <u>any</u> of the following:
 - a. Provider submits medical justification;
 - b. Documentation of recent (within the last 6 months) office visit or office chart notes demonstrating follow-up with the member;
 - c. Attestation that provider has reviewed the treatment plan with the member and assessed the risks and benefits of Nucynta ER;

4. If request is for a dose increase, new dose does not exceed 500 mg per day. Approval duration: 6 months or duration of request (whichever is less)

E. Other diagnoses/indications – Not applicable

III. Diagnoses/Indications for which coverage is NOT authorized - Not applicable

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym KeyFDA: Food and Drug AdministrationMME: morphine milligram equivalentsNSAID: non-steroidal anti-inflammatory
drugPA: prior authorizationREMS: Risk Evaluation and Mitigation
StrategySNRI: serotonin-norepinephrine reuptake
inhibitorTIRF: transmucosal immediate-release fentanyl
TCA: tricyclic antidepressant

Appendix B: Therapeutic Alternatives Not applicable

Page **5** of **9**



Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): significant respiratory depression; acute or severe bronchial asthma; gastrointestinal obstruction, including paralytic ileus; hypersensitivity to the opioid active ingredient, salts, or any component of the product; concurrent use of monoamine oxidase inhibitors or use of these within the last 14 days (Nucynta ER only).
- Boxed warning(s): potential for addiction, abuse, and misuse; Risk Evaluation and Mitigation Strategy (REMS); life-threatening respiratory depression; accidental ingestion; neonatal opioid withdrawal syndrome; cytochrome P450 3A4 interactions; risks from concomitant use with benzodiazepines or other CNS depressants.

Opioid Oral MME Conversion Factors		
Type of Opioid (strength units)	MME Conversion Factor	
Codeine (mg)	0.15	
Dihydrocodeine (mg)	0.25	
Fentanyl buccal or SL tablets, or lozenge/troche (mcg)	0.13	
Fentanyl film or oral spray (mcg)	0.18	
Fentanyl nasal spray (mcg)	0.16	
Fentanyl patch (mcg)	7.2	
Hydrocodone (mg)	1	
Hydromorphone (mg)	4	
Levorphanol tartrate (mg)	11	
Meperidine hydrochloride (mg)	0.1	
Methadone (mg)		
$>0, \le 20$	4	
$> 20, \le 40$	8	
$>40, \le 60$	10	
> 60	12	
Morphine (mg)	1	
Opium (mg)	1	
Oxycodone (mg)	1.5	
Oxymorphone (mg)	3	
Pentazocine (mg)	0.37	
Tapentadol (mg)	0.4	
Tramadol (mg)	0.1	

Appendix D: General Information

V. Dosage and Administration

There are numerous opioid analgesics, please refer to the package insert of your drug of interest for information on appropriate dosage and administration.



VI. Product Availability

There are numerous opioid analgesics, please refer to the package insert of your drug of interest for product availability information.

VII. References

- 1. Nucynta ER Prescribing Information. Stoughton, MA: Collegium Pharmaceutical, Inc.; December 2023. Available at: https://www.nucynta.com/. Accessed February 13, 2024.
- Dowell D, Ragan KR, Jones CM, Baldwin GT, and Chou R. CDC clinical practice guideline for prescribing opioids for pain – United States, 2022. MMWR Recomm Rep 2022;71(No. RR-3):1-95.
- 3. Kampman K, Jarvis M. American Society of Addiction Medicine (ASAM) national practice guideline for the use of medications in the treatment of addiction involving opioid use. J Addict Med. 2015 Sep-Oct;9(5):358-67.
- 4. Price R, Smith D, Franklin G, et al. Oral and topical treatment of painful diabetic polyneuropathy: Practice guideline update summary. Report of the AAN Guideline Subcommittee. Neurology 2022;98(1): 31-43.
- ElSayed NA, Aleppo G, Aroda VR, et al; on behalf of the American Diabetes Association, 12. Retinopathy, neuropathy, and foot Care: Standards of care in diabetes—2023. Diabetes Care 2023; 46 (Supplement 1): S203–S215.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2020 annual review; no significant changes; added HIM- Arkansas disclaimer re: coverage when the member has a terminal illness; references reviewed and updated.	11.26.19	02.20
Revised approval duration for short-acting agents from 30 days to 3 months, and for long-acting agents from 30 days to 12 months, per request from Ambetter Pharmacy Director and PA Ops.	12.01.20	12.20 (ad hoc)
1Q 2021 annual review: no significant changes; references reviewed and updated.	12.02.20	02.21
Revised wording in the Cancer/Palliative Care Initial Approval section to clarify the circumstances under which the requirement for a trial of two formulary short-acting agents would apply.	08.24.21	
1Q 2022 annual review: no significant changes; references reviewed and updated.	11.23.21	02.22
Removed reference to HIM.PA.103; aligned step therapy verbiage in sections IA,B,C with NF criteria in HIM.PA.103; For section IB (short-acting agents), removed "requiring PA" from title; For diabetic peripheral neuropathy, revised approval duration from 180 days to 6 months for initial therapy and 30 days to 6 months for continued therapy, removed drug specific (Nucynta ER) call out in title, and moved criteria to end of sections I and II.	07.18.22	



Reviews, Revisions, and Approvals	Date	P&T Approval Date
For section IC, clarified trial and failure language for short-acting and long-acting agent requests. Template changes applied to continued therapy section.	09.06.22	
Ad hoc update added Louisiana state specific regulations to section I.C.2.C.i-ii.	11.23.22	
1Q 2023 annual review: added sickle cell disease; for continued therapy for short-acting agents changed approval duration from 7 days to 1 month; references reviewed and updated.	12.01.22	02.23
For long acting agents or request exceeding 14-day supply within 28 days or 28-day supply within 90 days section, removed disclaimer for "if member is new to Centene benefit"; for section II.C.2 changed day supply requirement from "7-day supply of opioid in last 90 days" to "14-day supply of opioid within 28 days or 28-day supply within 90 days"; for section II.C II.D, added additional options to allow "documentation of recent (within the last 6 months) office visit or office chart notes demonstrating follow-up with the member" and "attestation that the provider has reviewed the treatment plan with the member and assessed the risks and benefits of the opioid dose and duration" to support need of opioid analgesics for member; for cancer, sickle cell disease, palliative care, removed Ohio specific 80 MME/day requirement.	03.02.23	
1Q 2024 annual review: for short-acting agents changed approval duration to 14 days; references reviewed and updated.	10.19.23	02.24
2Q 2024 annual review: no significant changes; references reviewed and updated.	01.19.24	05.24

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage



decisions and the administration of benefits are subject to all terms, conditions, exclusions, and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment, or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members, and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation.