

## Clinical Policy: Allogeneic Hematopoietic Cell Transplants for Sickle Cell Anemia and $\beta$ -Thalassemia

Reference Number: CP.MP.108

Last Review Date: 12/19

[Coding Implications](#)

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### Description

This policy describes the medical necessity requirements for allogeneic hematopoietic cell transplants for sickle cell anemia and  $\beta$ -thalassemia. Sickle cell anemia and  $\beta$ -thalassemia are two hemoglobinopathies caused by deleterious genetic alterations in hemoglobin. These monogenic diseases present a range of heterogeneous symptoms that stem from damaged red blood cell function. Despite their limitations, allogeneic hematopoietic cell transplants are the only curative therapies possible for these hemoglobinopathies.

### Policy/Criteria

- I. It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that allogeneic hematopoietic cell transplants for sickle cell anemia and homozygous  $\beta$ -thalassemia are **medically necessary** for members when all the following criteria are met:
  - A. *Sickle Cell Anemia*, meets all:
    1. Age  $\leq$  45 years (children and young adults);
    2. HLA-matched, first-degree relative donor is available;
    3. History of stroke or is at risk of stroke or end-organ damage, as shown by at least one of the following: prior stroke, recurrent acute chest syndrome, recurrent vaso-occlusive crises, or red blood cell alloimmunization on chronic transfusion therapy.
  - B. *Homozygous  $\beta$ -Thalassemia*, meets all:
    1. Age  $\leq$  45 years (children and young adults);
    2. HLA-matched donor is available, one of the following:
      - a. Cord blood is the source of stem cells, and the donor is a first-degree relative;
      - b. Bone marrow is the source of stem cells;
      - c. Peripheral blood is the source, and the donor is either unable to, or refuses to donate bone marrow;
    3. Transfusion-dependent due to thalassemia;
    4. A standard, myeloablative conditioning regimen will be used;
    5. Request is made by, or in consultation with, a specializing in treating thalassemia.
- II. It is the policy of health plans affiliated with Centene Corporation that the following are considered **experimental/investigational**:
  - A. Autologous hematopoietic cell transplant for sickle cell anemia;
  - B. Autologous hematopoietic cell transplant for  $\beta$ -thalassemia;
  - C. Allogeneic hematopoietic cell transplants for the treatment of sickle cell anemia or homozygous  $\beta$ -Thalassemia for any other indications than those specified above.

### Background

Hemoglobinopathies are a group of over 1,000 hematological disorders that result from deleterious molecular alterations to hemoglobin and are broadly classified into two categories

## CLINICAL POLICY

### Hematopoietic Cell Transplants for Sickle Cell Anemia and $\beta$ -Thalassemia

based on the phenotypic characteristics of these variations.<sup>1</sup> The first of these categories includes disorders, such as sickle cell anemia, in which there is a structural defect in one of the globin subunits.<sup>1</sup> Thalassemia belongs to the second category of hemoglobinopathies in which there is a quantitative defect in the production of one or more of the globin subunits.<sup>1</sup>

In adults, hemoglobin is a heterotetramer that is comprised of the  $\alpha$ - and  $\beta$ -globin subunits.<sup>2</sup> Each globin subunit forms a stable linkage with heme so that oxygen in the cytosol of an erythrocyte can bind reversibly to heme's iron atoms.<sup>2</sup> The hemoglobin tetramer  $\alpha_2\beta_2$  binds and unloads oxygen in a cooperative manner, which maximizes the transport of oxygen to cells.<sup>2</sup> Additional gas transport functions of hemoglobin include the transport of carbon dioxide and nitric oxide.<sup>3</sup> Each of these physiological aspects of hemoglobin are deleteriously affected in the hemoglobinopathy disorders.

#### *Sickle Cell Anemia and $\beta$ -Thalassemia*

Sickle cell disease results from a synonymous mutation that exchanges glutamic acid with valine at position 6 in the  $\beta$ -globin subunit.<sup>4</sup> Homozygous inheritance of this mutation results in the disease phenotype, whereas heterozygous carriers do not exhibit clinical disease symptoms; heterozygous carriers are also referred to as having sickle cell trait.<sup>4</sup> This amino acid substitution causes deoxygenated hemoglobin to rigid polymers in red blood cells, which ultimately forms the classic sickle-shaped morphology.<sup>2</sup> The sickle red blood cells occlude the microvasculature which leads to tissue hypoxia, infarction, and chronic hemolytic anemia.<sup>4</sup> Thus, sickle cell anemia presents a heterogeneous range of clinical manifestations, including pain, strokes, vaso-occlusive episodes, multi-organ injury, reduced quality of life, and shortened lifespan.<sup>2,4</sup>

Autosomal mutations in the gene encoding the  $\beta$ -globin subunit cause  $\beta$ -thalassemia (also known as thalassemia major or Cooley's anemia).<sup>5</sup> These mutations inhibit the synthesis of  $\beta$ -globin in erythropoietic cells.<sup>2,5</sup> The extent of the molecular basis for these mutations is very heterogeneous because over 200 mutations within the  $\beta$ -globin subunit, ranging from synonymous mutations to deletions.<sup>1</sup> Consequently,  $\alpha$ -globin molecules form toxic aggregates which destroy erythroid precursors through a process called ineffective erythropoiesis.<sup>2,5</sup> Also, individuals with  $\beta$ -thalassemia suffer from anemia due to shortened red blood cell survival, hemolysis anemia.<sup>5</sup>

#### *Hematopoietic Cell Transplantation*

Hematopoietic cell transplantation (HCT) is recognized as the only cure for sickle cell disease, and the success rate for specific pediatric groups has been shown to be 85 – 90%.<sup>4</sup> In the United States, it is estimated that the number of children with homozygous sickle cell anemia is 70,000 – 100,000, of which 5,000 – 7,000 could be eligible for transplantation.<sup>6</sup> A survey of the European Blood and Marrow Transplant and CIBMTR data files that ~1,200 patients in total have received HCT for sickle cell disease, and the 3 year survival rate is ~90% regardless of the source of hematopoietic stem cells.<sup>6</sup> Furthermore, Lucarelli *et al.* and Angelucci *et al.* both have documented the literature for recent reports on outcomes of HCT from HLA-matched donors in cases of  $\beta$ -thalassemia.<sup>8,9</sup> Although stem cell sources and the risk categories of the patients vary, overall survival and thalassemia-free survival range from approximately 65% - 90% among the numerous reports.<sup>8,9</sup>

**CLINICAL POLICY**

**Hematopoietic Cell Transplants for Sickle Cell Anemia and  $\beta$ -Thalassemia**

The establishment of complete donor-derived erythropoiesis can stabilize function in affected organs, such as the central nervous system and lungs.<sup>7</sup> However, HCT related organ toxicities, graft vs. host disease, graft rejection, and donor availability are major limitations of this procedure.<sup>7</sup> Infertility and gonadal failure are two specific morbidities with which HCT is associated.<sup>4</sup> Also, use of fully matched sibling donors as potentially eligible donors is one of the limitations for HCT implementation.<sup>4</sup> However, siblings are preferable HCT donors due to the lowered risk of graft vs host disease.<sup>6</sup>

Other differences between the considerations for HCT for  $\beta$ -thalassemia and sickle cell anemia include key issues for risk factors for transplant-related complications, transplant outcome, and conditioning regimen.<sup>9</sup> The major risk factors when considering HCT for  $\beta$ -thalassemia include age and organ dysfunction due to iron overload, whereas the major risk factors for HCT due to sickle cell anemia are age and history of cerebral events.<sup>9</sup> Control of iron overload and related tissue damage is a significant consideration for HCT for  $\beta$ -thalassemia, while obtaining a cure from chronic inflammation and prevention of sickle cell related organ damage must be considered for sickle cell anemia.<sup>9</sup> Lastly,  $\beta$ -thalassemia patients require an ablative conditioning regimen, whereas a reduced intensity regimen seems to induce stable chimerism and full donor erythropoiesis in sickle cell anemia patients.<sup>9</sup>

**Coding Implications**

This clinical policy references Current Procedural Terminology (CPT®). CPT® is a registered trademark of the American Medical Association. All CPT codes and descriptions are copyrighted 2019, American Medical Association. All rights reserved. CPT codes and CPT descriptions are from the current manuals and those included herein are not intended to be all-inclusive and are included for informational purposes only. The following codes are for informational purposes only. They are current at time of review of this policy. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

<b>CPT®*</b> <b>Codes</b>	<b>Description</b>
38205	Blood-derived hematopoietic progenitor cell harvesting for transplantation, per collection; allogeneic
38240	Hematopoietic progenitor cell (HPC); allogeneic transplantation per donor

<b>HCPCS</b> <b>Codes</b>	<b>Description</b>
S2150	Bone marrow or blood-derived stem cells (peripheral or umbilical), allogeneic or autologous, harvesting, transplantation, and related complications; including: pheresis and cell preparation/storage; marrow ablative therapy; drugs, supplies, hospitalization with outpatient follow-up; medical/surgical, diagnostic, emergency, and rehabilitative services; and the number of days of pre and post-transplant care in the global definition

Hematopoietic Cell Transplants for Sickle Cell Anemia and  $\beta$ -Thalassemia

ICD-10-CM Code	Description
D56.1	Beta thalassemia
D57.0-D57.819	Sickle-cell disorders

Reviews, Revisions, and Approvals	Date	Approval Date
Policy developed	03/16	03/16
References reviewed and updated	03/17	03/17
Changed age restriction for sickle cell from $\leq 25$ to $\leq 45$ , and changed age for beta thalassemia from $\leq 30$ to $\leq 45$ . Revised I.B.3 to require transfusion dependence due to thalassemia and removed “deterioration with conventional treatments...” Added I.B.4 Request is made by, or in consultation with, a thalassemia specialist	11/17	11/17
Clarified language in II.C. References reviewed and updated.	02/18	02/18
Sickle cell: specified that donor should be a first-degree relative, and that the conditioning regimen should be myeloablative. Beta thalassemia: added that cord blood is allowed if donated by a first-degree relative, added bone marrow as an acceptable source, and peripheral blood as an acceptable source if the donor is unable or unwilling to donate bone marrow; changed requirement for thalassemia specialist to “provider specializing in thalassemia”	12/18	01/19
Removed I.A. 4. Requirement of a standard, myeloablative conditioning regimen. References reviewed and updated. Specialist review.	11/19	12/19

**References**

1. Forget BG, Bunn HF. "Classification of the Disorders of Hemoglobin." *Cold Spring Harbor perspectives in medicine* 3.2 (2013): a011684.
2. Sankaran, Vijay G., and Mitchell J. Weiss. "Anemia: progress in molecular mechanisms and therapies." *Nature medicine* 21.3 (2015): 221-230.
3. Schechter, Alan N. "Hemoglobin research and the origins of molecular medicine." *Blood* 112.10 (2008): 3927-3938.
4. Kanter, Julie, and Rebecca Kruse-Jarres. "Management of sickle cell disease from childhood through adulthood." *Blood reviews* 27.6 (2013): 279-287.
5. Ribeil, Jean-Antoine, et al. "Ineffective erythropoiesis in  $\beta$ -thalassemia." *The Scientific World Journal* 2013 (2013).
6. Gluckman, Eliane. "Allogeneic transplantation strategies including haploidentical transplantation in sickle cell disease." *ASH Education Program Book 2013.1* (2013): 370-376.
7. Shenoy, Shalini. "Hematopoietic stem-cell transplantation for sickle cell disease: current evidence and opinions." *Therapeutic advances in hematology* 4.5 (2013): 335-344.
8. Lucarelli, Guido, et al. "Hematopoietic stem cell transplantation in thalassemia and sickle cell anemia." *Cold Spring Harbor perspectives in medicine* 2.5 (2012): a011825.
9. Angelucci, Emanuele, et al. "Hematopoietic stem cell transplantation in thalassemia major and sickle cell disease: indications and management recommendations from an international expert panel." *Haematologica* 99.5 (2014): 811-820.

## CLINICAL POLICY

### Hematopoietic Cell Transplants for Sickle Cell Anemia and $\beta$ -Thalassemia

10. Khan S, Rodgers GP. Hematopoietic cell transplantation in sickle cell disease. In: UpToDate, Mahoney DH, Chao NJ. (Eds), UpToDate, Waltham, MA. Accessed Nov 14, 2019
11. Hayes Health Technology. Allogeneic Hematopoietic Stem Cell Transplantation for Sickle Cell Disease in Children. Lansdale, PA: Hayes, Inc.; 2014 Winifred S. Hayes, Inc. February 25, 2016. Update March 20, 2019. Accessed Nov 15, 2019.
12. Ozdogu H, Boga C, Yeral M, et al. Allogeneic peripheral stem cell transplantation from HLA-matched related donors for adult sickle cell disease: remarkable outcomes from a single-center trial. *Bone Marrow Transplant.* 2018 Feb 5.
13. Gilman AL, Eckrich MJ, Epstein S, et al. Alternative donor hematopoietic stem cell transplantation for sickle cell disease. *Blood Adv.* 2017 Jun 28;1(16):1215-1223.
14. Allen ES, Srivastava K, Hsieh MM, et al. Immunohaematological complications in patients with sickle cell disease after haemopoietic progenitor cell transplantation: a prospective, single-centre, observational study. *Lancet Haematol.* 2017 Nov;4(11):e553-e561
15. Rangarajan HG, Abu-Arja R, Pai V, Guilcher GMT, Soni S. Outcomes of Unrelated Donor Stem Cell Transplantation with Post-Transplant Cyclophosphamide for Graft-versus-Host Disease Prophylaxis in Patients with Severe Sickle Cell Disease. *Biol Blood Marrow Transplant.* 2018 Feb;24(2):413-417.
16. Benz EJ, Angelucci E. Management and prognosis of the thalassemias. In: UpToDate, Vichinsky EP (Ed), UpToDate, Waltham, MA. Accessed Dec 18, 2018.
17. Angelucci E, Benz EJ. Hematopoietic cell transplantation for transfusion-dependent thalassemia. In: UpToDate, Negrin RS (Ed), UpToDate, Waltham, MA. Accessed Nov 15, 2019.
19. Ghavamzadeh A, Kasaeian A, Rostami T, Kiumarsi A. Comparable outcomes of allogeneic peripheral blood versus bone marrow hematopoietic stem cell transplantation in major thalassemia: a multivariate long-term cohort analysis. *Biol Blood Marrow Transplant.* 2018 Sep 26. pii: S1083-8791(18)30593-7. doi: 10.1016/j.bbmt.2018.09.026.

#### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and

## CLINICAL POLICY

### Hematopoietic Cell Transplants for Sickle Cell Anemia and $\beta$ -Thalassemia

limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

**Note: For Medicaid members**, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

**Note: For Medicare members**, to ensure consistency with the Medicare National Coverage Determinations (NCD) and Local Coverage Determinations (LCD), all applicable NCDs, LCDs, and Medicare Coverage Articles should be reviewed prior to applying the criteria set forth in this clinical policy. Refer to the CMS website at <http://www.cms.gov> for additional information.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or

**CLINICAL POLICY**

**Hematopoietic Cell Transplants for Sickle Cell Anemia and  $\beta$ -Thalassemia**

remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.