

Clinical Policy: Benralizumab (Fasenra)

Reference Number: CP.PHAR.373

Effective Date: 06.01.18 Last Review Date: 02.20

Line of Business: Commercial, HIM, Medicaid

Coding Implications
Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Benralizumab (FasenraTM) is an interleukin (IL)-5 receptor alpha-directed cytolytic monoclonal antibody.

FDA Approved Indication(s)

Fasenra is indicated for the add-on maintenance treatment of patients with severe asthma aged 12 years and older, and with an eosinophilic phenotype.

Limitation(s) of use:

- Fasenra is not indicated for treatment of other eosinophilic conditions.
- Fasenra is not indicated for the relief of acute bronchospasm or status asthmaticus.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation® that Fasenra is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Severe Asthma (must meet all):
 - 1. Diagnosis of asthma;
 - 2. Member has an absolute blood eosinophil count ≥ 150 cells/mcL within the past 3 months:
 - 3. Prescribed by or in consultation with a pulmonologist, immunologist, or allergist;
 - 4. Age \geq 12 years;
 - 5. Member has experienced ≥ 2 exacerbations within the last 12 months, requiring any of the following despite adherent use of controller therapy (i.e., medium- to high-dose inhaled corticosteroid (ICS) plus either a long-acting beta₂ agonist (LABA) or leukotriene modifier (LTRA) if LABA contraindication/intolerance):
 - a. Oral/systemic corticosteroid treatment (or increase in dose if already on oral corticosteroid);
 - b. Urgent care visit or hospital admission;
 - c. Intubation;
 - 6. Fasenra is prescribed concurrently with an ICS plus either a LABA or LTRA;
 - 7. Fasenra is not prescribed concurrently with Cinqair®, Nucala®, Dupixent®, or Xolair®;



8. Dose does not exceed 30 mg every 4 weeks for the first 3 doses, then 30 mg every 8 weeks thereafter.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Severe Asthma (must meet all):

- 1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
- 2. Demonstrated adherence to asthma controller therapy that includes an ICS plus either a LABA or LTRA;
- 3. Member is responding positively to therapy (examples may include but are not limited to: reduction in exacerbations or corticosteroid dose, improvement in forced expiratory volume over one second since baseline, reduction in the use of rescue therapy);
- 4. Fasenra is not prescribed concurrently with Cinqair, Nucala, Dupixent, or Xolair;
- 5. If request is for a dose increase, new dose does not exceed 30 mg every 8 weeks.

Approval duration:

Medicaid/HIM – 12 months

Commercial – 6 months or member's renewal period, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

- 1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
 - Approval duration: Duration of request or 6 months (whichever is less); or
- Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- **A.** Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies CP.CPA.09 for commercial, HIM.PHAR.21 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents;
- **B.** Acute bronchospasm or status asthmaticus.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

FDA: Food and Drug Administration GINA: Global Initiative for Asthma

BEC: blood eosinophil count ICS: inhaled corticosteroid



IL: interleukin LTRA: leukotriene modifier

LABA: long-acting beta₂ agonist

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
ICS (medium – high dose		77203333
Qvar® (beclomethasone)	> 200 mcg/day 40 mcg, 80 mcg per actuation 1-4 actuations BID	4 actuations BID
budesonide (Pulmicort®)	> 400 mcg/day 90 mcg, 180 mcg per actuation 2-4 actuations BID	2 actuations BID
Alvesco® (ciclesonide)	> 160 mcg/day 80 mcg, 160 mcg per actuation 1-2 actuations BID	2 actuations BID
Aerospan® (flunisolide)	> 320 mcg/day 80 mcg per actuation 2-4 actuations BID	2 actuations BID
Flovent® (fluticasone propionate)	> 250 mcg/day 44-250 mcg per actuation 2-4 actuations BID	2 actuations BID
Arnuity Ellipta® (fluticasone furoate)	200 mcg/day 100 mcg, 200 mcg per actuation 1 actuation QD	1 actuation QD
Asmanex® (mometasone)	>220 mcg/day HFA: 100 mcg, 200 mcg per actuation Twisthaler: 110 mcg, 220 mcg per actuation 1-2 actuations QD to BID	2 inhalations BID
LABA		
Serevent® (salmeterol)	50 mcg per dose 1 inhalation BID	1 inhalation BID
Combination products (I		
Dulera® (mometasone/ formoterol)	100/5 mcg, 200/5 mcg per actuation 2 actuations BID	4 actuations per day
Breo Ellipta® (fluticasone/vilanterol)	100/25 mcg, 200/25 mcg per actuation 1 actuation QD	1 actuation QD
Advair® (fluticasone/ salmeterol)	Diskus: 100/50 mcg, 250/50 mcg, 500/50 mcg per actuation HFA: 45/21 mcg, 115/21 mcg, 230/21 mcg per actuation 1 actuation BID	1 actuation BID



Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose	
fluticasone/salmeterol	55/13 mcg, 113/14 mcg, 232/14 mcg	1 actuation BID	
(Airduo RespiClick®)	per actuation		
	1 actuation BID		
Symbicort® (budesonide/	80 mcg/4.5 mcg, 160 mcg/4.5 mcg per	2 actuations BID	
formoterol)	actuation		
,	2 actuations BID		
LTRA			
montelukast (Singulair®)	4 to 10 mg PO QD	10 mg per day	
zafirlukast (Accolate®)	10 to 20 mg PO BID	40 mg per day	
zileuton ER (Zyflo® CR)	1200 mg PO BID	2,400 mg per day	
Zyflo® (zileuton)	600 mg PO QID	2,400 mg per day	
Oral corticosteroids			
dexamethasone	0.75 to 9 mg/day PO in 2 to 4 divided	Varies	
(Decadron®)	doses		
methylprednisolone	40 to 80 mg PO in 1 to 2 divided doses	Varies	
(Medrol®)	_		
prednisolone (Millipred®,	40 to 80 mg PO in 1 to 2 divided doses	Varies	
Orapred ODT®)			
prednisone (Deltasone®)	40 to 80 mg PO in 1 to 2 divided doses	Varies	

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

• Contraindication(s): hypersensitivity

• Boxed warning(s): none reported

Appendix D: General Information

- The pivotal trials defined severe asthma as 2 or more exacerbations of asthma despite regular use of high-dose ICS plus an additional controller (e.g., LABA or LTRA) with or without oral corticosteroids. Although the CALIMA trial included patients receiving medium-dose ICS, Fasenra was not shown to have an effect on annual exacerbation rate, pre-bronchodilator forced expiratory volume in 1 second, or total asthma symptom score in those patients.
- Clinically significant exacerbation was defined as a worsening of asthma (any new or increased symptoms or signs that were concerning) that led to one of the following: (1) use of systemic corticosteroids, (2) emergency department or visit to urgent care center, or (3) inpatient hospital stay.
- Baseline blood eosinophil count (BEC) is a predictor of response to therapy. Although
 the SIROCCO and CALIMA trials were powered for efficacy analysis in patients with
 baseline BEC ≥ 300 cells/μL, a pooled analysis which stratified patients by baseline BEC
 (≥ 0 cells/μL, ≥ 150 cells/μL, ≥ 300 cells/μL, and ≥ 450 cells/μL) found Fasenra to have
 a statistically significant positive treatment effect on those with baseline BEC ≥ 150



- cells/ μ L. In addition, the ZONDA trial found Fasenra to significantly reduce oral corticosteroid dose in patients with baseline BEC \geq 150 cells/ μ L.
- Patients could potentially meet asthma criteria for both Xolair and Fasenra. The
 combination has not been studied. Approximately 30% of patients in the Nucala MENSA
 study also were candidates for therapy with Xolair.
- Lab results for blood eosinophil counts can be converted into cells/mcL using the following unit conversion calculator: https://www.fasenrahcp.com/m/fasenra-eosinophil-calculator.html.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Severe asthma	30 mg SC every 4 weeks for the first 3 doses,	See regimen
	followed by once every 8 weeks thereafter	

VI. Product Availability

- Single-dose prefilled syringe with solution for injection: 30 mg/mL
- Single-dose autoinjector Fasenra Pen with solution for injection: 30 mg/mL

VII. References

- 1. Fasenra Prescribing Information. Wilmington, DE: AstraZeneca Pharmaceuticals LP; October 2019. Available at: www.fasenra.com. Accessed November 7, 2019.
- 2. National Asthma Education and Prevention Program: Expert panel report III: Guidelines for the diagnosis and management of asthma. Bethesda, MD: National Heart, Lung, and Blood Institute, 2007. (NIH publication no. 08-4051). Available at http://www.nhlbi.nih.gov/health-pro/guidelines/current/asthma-guidelines. Accessed November 7, 2019.
- 3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2019. Available at: http://www.clinicalpharmacology.com. Accessed November 7, 2019.
- 4. Global Initiative for Asthma. Global strategy for asthma management and prevention, 2019. Available at: www.ginasthma.org. Published June 2019. Accessed November 5, 2019.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
C9466	Injection, benralizumab, 1 mg

Reviews, Revisions, and Approvals	Date	P&T
		Approval Date
Policy created	01.16.18	05.18
1Q 2019 annual review: modified ICS requirement to include	10.11.18	02.19
medium dose ICS per GINA 2018 recommendations; added option		
for immunologist prescribing; link to blood eosinophil unit		



Reviews, Revisions, and Approvals	Date	P&T Approval Date
conversion calculator added to Appendix D; references reviewed		
and updated.		
Added HIM-Medical Benefit line of business.	05.22.19	
1Q 2020 annual review: revised HIM-Medical Benefit to HIM line	11.07.19	02.20
of business; added requirement that Fasenra is not prescribed		
concurrently with other biologic therapies for asthma; added new		
autoinjector formulation; references reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.



Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation.

Corporation are registered trademarks exclusively owned by Centene Corporation.