

Clinical Policy: Sargramostim (Leukine)

Reference Number: CP.PHAR.295

Effective Date: 12.01.16

Last Review Date: 08.22

Line of Business: Commercial, HIM, Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sargramostim (Leukine[®]) is a recombinant human granulocyte-macrophage colony stimulating factor (GM-CSF).

FDA Approved Indication(s)

Leukine is indicated:

- To shorten time to neutrophil recovery and to reduce the incidence of severe and life-threatening infections and infections resulting in death following induction chemotherapy in adult patients 55 years and older with acute myeloid leukemia (AML);
- For the mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation in adult patients;
- For the acceleration of myeloid reconstitution following autologous peripheral blood progenitor cell (PBPC) or bone marrow transplantation in adult and pediatric patients 2 years of age and older with non-Hodgkin's lymphoma (NHL), acute lymphoblastic leukemia (ALL) and Hodgkin's lymphoma (HL);
- For the acceleration of myeloid reconstitution following allogeneic bone marrow transplantation in adult and pediatric patients 2 years of age and older;
- For treatment of delayed neutrophil recovery or graft failure after autologous or allogeneic bone marrow transplantation in adult and pediatric patients 2 years of age and older;
- To increase survival in adult and pediatric patients from birth to 17 years of age acutely exposed to myelosuppressive doses of radiation (Hematopoietic Syndrome of Acute Radiation Syndrome [H-ARS]).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Leukine is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Acute Myelogenous Leukemia (must meet all):**

1. Diagnosis of AML;
2. Prescribed for use following induction therapy for AML;
3. Age \geq 55 years;

4. Member meets one of the following (a or b):
 - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
 - b. Failure of Zarxio[®] unless contraindicated or clinically significant adverse effects are experienced;
**Prior authorization may be required for Zarxio*
5. Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle;
6. Dose does not exceed 250 mcg/m² IV daily.

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

B. Peripheral Blood Progenitor Cell Collection and Transplantation (must meet all):

1. Prescribed for one of the following (a or b):
 - a. Mobilization of hematopoietic progenitor cells into peripheral blood for collection by leukapheresis and autologous transplantation;
 - b. Following autologous PBPC transplantation in members with NHL, ALL, HL for acceleration of myeloid reconstitution;
2. Age ≥ 2 years;
3. Member meets one of the following (a or b):
 - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
 - b. Failure of Zarxio[®] unless contraindicated or clinically significant adverse effects are experienced;
**Prior authorization may be required for Zarxio*
4. Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle;
5. Dose does not exceed 250 mcg/m² IV or SC daily.

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

C. Bone Marrow Transplantation (must meet all):

1. Prescribed for use in one of the following settings (a, b, or c):
 - a. Following autologous BMT in members with NHL, ALL, or HL for acceleration of myeloid reconstitution;
 - b. Following allogeneic BMT for acceleration of myeloid reconstitution;
 - c. Following BMT where engraftment is delayed or has failed;
2. Age ≥ 2 years;
3. Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle;
4. Member meets one of the following (a or b):
 - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
 - b. Failure of Zarxio[®] unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Zarxio*

5. Dose does not exceed 500 mcg/m² IV daily.

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

D. Acute Radiation Syndrome (must meet all):

1. Prescribed for use following suspected or confirmed acute exposure to myelosuppressive doses of radiation;
2. Member meets one of the following (a or b):
 - a. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
 - b. Failure of Zarxio[®] unless contraindicated or clinically significant adverse effects are experienced;

**Prior authorization may be required for Zarxio*

3. Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle;
4. Dose does not exceed one of the following (a, b, or c):
 - a. Weight <15 kg: 12 mcg/kg SC daily;
 - b. Weight 15 kg to 40 kg: 10 mcg/kg SC daily;
 - c. Weight > 40 kg: 7 mcg/kg SC daily.

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

E. Other diagnoses/indications (must meet 1 and 2)

1. Member meets one of the following (a or b):
 - a. Member must use Zarxio, unless both are contraindicated or clinically significant adverse effects are experienced;*
2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

**Prior authorization may be required for Zarxio*

II. Continued Therapy

A. All Indications in Section I (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy;
3. Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle;
4. Dose does not exceed the FDA-approved maximum recommended dose for the relevant indication in Section V.

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 and 2):

1. Member meets one of the following (a or b):
 - a. Member must use Zarxio, unless both are contraindicated or clinically significant adverse effects are experienced;*
*Prior authorization may be required for Zarxio
 - b. Request is for treatment associated with cancer for a State with regulations against step therapy in certain oncology settings (*see Appendix E*);
2. One of the following (a or b):
 - a. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.
Approval duration: Duration of request or 6 months (whichever is less); or
 - b. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

ALL: acute lymphoblastic leukemia
AML: acute myelogenous leukemia
BMT: bone marrow transplantation
FDA: Food and Drug Administration
GM-CSF: granulocyte-macrophage colony stimulating factor

H-ARS: hematopoietic syndrome of acute radiation syndrome
NHL: non-Hodgkin's lymphoma
PBPC: peripheral blood progenitor cell

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent and may require prior authorization.

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
Zarxio (filgrastim-sndz),	AML: 5 mcg/kg SC or IV QD BMT: 10 mcg/kg IV or SC infusion QD	AML: 30 mcg/kg/day [IV] or 24 mcg/kg/day [SC] BMT,

Drug	Dosing Regimen	Dose Limit/ Maximum Dose
	PBPC collection: 10 mcg/kg SC bolus or continuous infusion QD Acute Radiation Syndrome: 10 mcg/kg SC QD	PBPC collection, Acute Radiation Syndrome: 10 mcg/kg/day

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): history of serious allergic reactions, including anaphylaxis
- Boxed warning(s): none reported

Appendix D: General Information

- Because of potential sensitivity of rapidly dividing hematopoietic progenitor cells, Leukine should not be administered simultaneously with cytotoxic chemotherapy or radiotherapy or within 24 hours preceding or following chemotherapy or radiotherapy.
- Use Leukine with caution in patients with pre-existing fluid retention, pulmonary infiltrates, or congestive heart failure.

Appendix E: States with Regulations against Redirections in Cancer

State	Step Therapy Prohibited?	Notes
FL	Yes	For stage 4 metastatic cancer and associated conditions.
GA	Yes	For stage 4 metastatic cancer. Redirection does not refer to review of medical necessity or clinical appropriateness.
IA	Yes	For standard of care stage 4 cancer drug use, supported by peer-reviewed, evidence-based literature, and approved by FDA.
LA	Yes	For stage 4 advanced, metastatic cancer or associated conditions. Exception if “clinically equivalent therapy, contains identical active ingredient(s), and proven to have same efficacy.
NV	Yes	Stage 3 and stage 4 cancer patients for a prescription drug to treat the cancer or any symptom thereof of the covered person
OH	Yes	<i>*Applies to Commercial and HIM requests only*</i> For stage 4 metastatic cancer and associated conditions
PA	Yes	For stage 4 advanced, metastatic cancer
TN	Yes	For advanced metastatic cancer and associated conditions
TX	Yes	For stage 4 advanced, metastatic cancer and associated conditions

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
AML	250 mcg/m ² /day IV over a 4 hour period approximately on day 11 or	250 mcg/m ² IV daily

Indication	Dosing Regimen	Maximum Dose
	four days following the completion of induction chemotherapy	
PBPC collection and transplantation	250 mcg/m ² /day administered IV over 24 hours or SC once daily	250 mcg/m ² IV or SC daily
Myeloid reconstitution after autologous or allogeneic BMT	250 mcg/m ² /day IV over a 2 hour period beginning two to four hours after bone marrow infusion, and not less than 24 hours after the last dose of chemotherapy or radiotherapy	500 mcg/m ² IV daily
BMT failure or engraftment delay	250 mcg/m ² /day for 14 days as a 2 hour IV infusion	500 mcg/m ² IV daily
Acute Radiation Syndrome	Weight-based dose SC QD: >40 kg: 7 mcg/kg 15 to 40 kg: 10 mcg/kg <15 kg: 12 mcg/kg	See dosing regimen

VI. Product Availability

Lyophilized powder: 250 mcg single-dose vial
Solution: 500 mcg/mL multiple-dose vial

VII. References

1. Leukine Prescribing Information. Bridgewater, NJ: Sanofi-Aventis U.S., LLC.; March 2018. Available at: www.leukine.com. Accessed April 20, 2022.
2. National Comprehensive Cancer Network: Hematopoietic Growth Factors Version 1.2022. Available at: https://www.nccn.org/professionals/physician_gls/pdf/growthfactors.pdf. Accessed: April 20, 2022.
3. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed April 20, 2022.
4. DRUGDEX[®] System [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed April 20, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPSC Codes	Description
J2820	Injection, sargramostim (GM-CSF), 50 mcg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
3Q 2018 annual review: added HIM line of business, added new indication for acute radiation syndrome; removed contraindications	05.02.18	08.18

Reviews, Revisions, and Approvals	Date	P&T Approval Date
that are no longer included in the product label; modified age restrictions consistent with label; references reviewed and updated.		
3Q 2019 annual review: no significant changes; references reviewed and updated.	05.15.19	08.19
3Q 2020 annual review: added Commercial line of business (retire CP.CPA.262); for ARS indication added weight based dosing to criteria set; references reviewed and updated.	05.04.20	08.20
Added redirection to Zarxio for all indications per prior clinical guidance and data analytics financial analysis.	10.12.20	
3Q 2021 annual review: no significant changes; allowed by-passing of redirection if state regulations do not allow step therapy in Stage IV or metastatic cancer settings for AML; modified HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	04.05.21	08.21
Added Nevada to Appendix E.	08.03.21	
3Q 2022 annual review: removed general description of “stage IV or metastatic” cancer for states with regulations against redirections; applied redirection bypass for State with regulations against step therapy to all indications; added requirement that Leukine will not be prescribed concurrently with other colony stimulating factors (e.g., filgrastim, pegfilgrastim) within any chemotherapy cycle; references reviewed and updated.	04.20.22	08.22

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note: For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2016 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.