

Clinical Policy: Sarilumab (Kevzara)

Reference Number: CP.PHAR.346

Effective Date: 07.18.17

Last Review Date: 02.21

Line of Business: Medicaid

[Coding Implications](#)[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

Description

Sarilumab (Kevzara[®]) is an interleukin-6 (IL-6) receptor antagonist.

FDA Approved Indication(s)

Kevzara is indicated for treatment of adult patients with moderately to severely active rheumatoid arthritis (RA) who have had an inadequate response or intolerance to one or more disease-modifying antirheumatic drugs (DMARDs).

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Kevzara is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria**A. Rheumatoid Arthritis** (must meet all):

1. Diagnosis of RA per American College of Rheumatology (ACR) criteria (*see Appendix E*);
2. Prescribed by or in consultation with a rheumatologist;
3. Age \geq 18 years;
4. Member meets one of the following (a or b):
 - a. Failure of a \geq 3 consecutive month trial of methotrexate (MTX) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effect are experienced;
 - b. If intolerance or contraindication to MTX (*see Appendix D*), failure of a \geq 3 consecutive month trial of at least ONE conventional DMARD (e.g., sulfasalazine, leflunomide, hydroxychloroquine) at up to maximally indicated doses, unless contraindicated or clinically significant adverse effect are experienced;
5. Documentation of one of the following baseline assessment scores (a or b):
 - a. Clinical disease activity index (CDAI) score (*see Appendix F*);
 - b. Routine assessment of patient index data 3 (RAPID3) score (*see Appendix G*);
6. Dose does not exceed 200 mg every two weeks.

Approval duration: 6 months

B. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Rheumatoid Arthritis (must meet all):

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
2. Member is responding positively to therapy as evidenced by one of the following (a or b):
 - a. A decrease in CDAI (*see Appendix F*) or RAPID3 (*see Appendix G*) score from baseline;
 - b. Medical justification stating inability to conduct CDAI re-assessment, and submission of RAPID3 score associated with disease severity that is similar to initial CDAI assessment or improved;
3. If request is for a dose increase, new dose does not exceed 200 mg every two weeks.

Approval duration: 12 months

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – CP.PMN.53 for Medicaid or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

CDAI: clinical disease activity index

DMARD: disease-modifying
antirheumatic drug

FDA: Food and Drug Administration

IL-6: interleukin-6

MTX: methotrexate

RA: rheumatoid arthritis

RAPID3: routine assessment of patient
index data 3

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
azathioprine (Azasan [®] , Imuran [®])	RA 1 mg/kg/day PO QD or divided BID	2.5 mg/kg/day
Cuprimine [®] (d-penicillamine)	RA* <u>Initial dose:</u> 125 or 250 mg PO QD <u>Maintenance dose:</u> 500 – 750 mg/day PO QD	1,500 mg/day
cyclosporine (Sandimmune [®] , Neoral [®])	RA 2.5 – 4 mg/kg/day PO divided BID	4 mg/kg/day
hydroxychloroquine (Plaquenil [®])	RA* <u>Initial dose:</u> 400 – 600 mg/day PO QD <u>Maintenance dose:</u> 200 – 400 mg/day PO QD	600 mg/day
leflunomide (Arava [®])	RA 100 mg PO QD for 3 days, then 20 mg PO QD	20 mg/day
methotrexate (Rheumatrex [®])	RA 7.5 mg/week PO, SC, or IM or 2.5 mg PO Q12 hr for 3 doses/week	30 mg/week
Ridaura [®] (auranofin)	RA 6 mg PO QD or 3 mg PO BID	9 mg/day (3 mg TID)
sulfasalazine (Azulfidine [®])	RA 2 g/day PO in divided doses	3 g/day

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

**Off-label*

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): known hypersensitivity to sarilumab or any of the inactive ingredients
- Boxed warning(s): risk of serious infections

Appendix D: General Information

- Definition of MTX or DMARD Failure
 - Child-bearing age is not considered a contraindication for use of MTX. Each drug has risks in pregnancy. An educated patient and family planning would allow use of MTX in patients who have no intention of immediate pregnancy.
 - Social use of alcohol is not considered a contraindication for use of MTX. MTX may only be contraindicated if patients choose to drink over 14 units of alcohol per week. However, excessive alcohol drinking can lead to worsening of the condition, so

- patients who are serious about clinical response to therapy should refrain from excessive alcohol consumption.
- Examples of positive response to therapy may include, but are not limited to:
 - Reduction in joint pain/swelling/tenderness
 - Improvement in ESR/CRP levels
 - Improvements in activities of daily living

Appendix E: The 2010 ACR Classification Criteria for RA

Add score of categories A through D; a score of ≥ 6 out of 10 is needed for classification of a patient as having definite RA.

A	Joint involvement	Score
	1 large joint	0
	2-10 large joints	1
	1-3 small joints (with or without involvement of large joints)	2
	4-10 small joints (with or without involvement of large joints)	3
	> 10 joints (at least one small joint)	5
B	Serology (at least one test result is needed for classification)	
	Negative rheumatoid factor (RF) <i>and</i> negative anti-citrullinated protein antibody (ACPA)	0
	Low positive RF <i>or</i> low positive ACPA <i>* Low: $< 3 \times$ upper limit of normal</i>	2
	High positive RF <i>or</i> high positive ACPA <i>* High: $\geq 3 \times$ upper limit of normal</i>	3
C	Acute phase reactants (at least one test result is needed for classification)	
	Normal C-reactive protein (CRP) and normal erythrocyte sedimentation rate (ESR)	0
	Abnormal CRP or abnormal ESR	1
D	Duration of symptoms	
	< 6 weeks	0
	≥ 6 weeks	1

Appendix F: Clinical Disease Activity Index (CDAI) Score

The Clinical Disease Activity Index (CDAI) is a composite index for assessing disease activity in RA. CDAI is based on the simple summation of the count of swollen/tender joint count of 28 joints along with patient and physician global assessment on VAS (0–10 cm) Scale for estimating disease activity. The CDAI score ranges from 0 to 76.

CDAI Score	Disease state interpretation
≤ 2.8	Remission
2.8 to ≤ 10	Low disease activity
10 to ≤ 22	Moderate disease activity
> 22	High disease activity

Appendix G: Routine Assessment of Patient Index Data 3 (RAPID3) Score

The Routine Assessment of Patient Index Data 3 (RAPID3) is a pooled index of the three patient-reported ACR core data set measures: function, pain, and patient global estimate of

status. Each of the individual measures is scored 0 – 10, and the maximum achievable score is 30.

RAPID3 Score	Disease state interpretation
≤ 3	Remission
3.1 to 6	Low disease activity
6.1 to 12	Moderate disease activity
> 12	High disease activity

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
RA	200 mg SC once every two weeks	200 mg every 2 weeks

VI. Product Availability

Single-dose prefilled syringe/pen: 150 mg/1.14 mL, 200 mg/1.14 mL

VII. References

1. Kevzara Prescribing Information. Bridgewater, NJ: Sanofi-Aventis U.S. LLC; April 2018. Available at: <https://www.kevzara.com/>. Accessed February 26, 2020.
2. Singh JA., Saag KG, Bridges SL, et al. 2015 American College of Rheumatology guideline for the treatment of rheumatoid arthritis. *Arthritis Care & Research*, 68: 1–25. doi:10.1002/acr.22783.
3. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2020. Available at: <http://www.clinicalpharmacology-ip.com/>. Accessed February 26, 2020.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-to-date sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J3590	Unclassified biologics

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	06.17	11.17
2Q 2018 annual review: removed TB testing requirement; references reviewed and updated.	02.27.18	05.18
4Q 2018 annual review: no significant changes; references reviewed and updated.	09.04.18	11.18
2Q 2019 annual review: no significant changes; added HIM-Medical Benefit; references reviewed and updated.	02.26.19	05.19
Removed HIM-Medical Benefit line of business; updated preferred redirections based on SDC recommendations and prior clinical guidance: for RA, removed trial of etanercept and adalimumab.	12.16.19	

Reviews, Revisions, and Approvals	Date	P&T Approval Date
2Q 2020 annual review: for RA, added specific diagnostic criteria for definite RA, baseline CDAI score requirement, and decrease in CDAI score as positive response to therapy; references reviewed and updated.	04.23.20	05.20
Revised typo in Appendix E from “normal ESR” to “abnormal ESR” for a point gained for ACR Classification Criteria.	11.22.20	
Added criteria for RAPID3 assessment for RA given limited in-person visits during COVID-19 pandemic, updated appendices; added coding implications,	11.24.20	02.21

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.