

Clinical Policy: Cemiplimab-rwlc (Libtayo)

Reference Number: CP.PHAR.397 Effective Date: 10.16.18 Last Review Date: 11.22 Line of Business: Commercial, HIM, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Cemiplimab-rwlc (Libtayo[®]) is a programmed death receptor-1 (PD-1) blocking antibody.

FDA Approved Indication(s)

Libtayo is indicated:

- For the treatment of patients with metastatic cutaneous squamous cell carcinoma (CSCC) or locally advanced CSCC who are not candidates for curative surgery or curative radiation.
- For the treatment of patients with locally advanced basal cell carcinoma (BCC) previously treated with a hedgehog pathway inhibitor or for whom a hedgehog pathway inhibitor is not appropriate.
- For the treatment of patients with metastatic BCC previously treated with a hedgehog pathway inhibitor or for whom a hedgehog pathway inhibitor is not appropriate.*
- In combination with platinum-based chemotherapy for the first-line treatment of adult patients with non-small cell lung cancer (NSCLC) with no epidermal growth factor receptor (EGFR), anaplastic lymphoma kinase (ALK) or ROS1 aberrations and is locally advanced where patients are not candidates for surgical resection or definitive chemoradiation or metastatic.
- As a single agent for the first-line treatment of patients with NSCLC whose tumors have high PD-L1 expression [Tumor Proportion Score (TPS) ≥ 50%] as determined by an FDA-approved test, with no EGFR, ALK or ROS1 aberrations, and is locally advanced where patients are not candidates for surgical resection or definitive chemoradiation or metastatic.

*The metastatic BCC indication is approved under accelerated approval based on tumor response rate and durability of response. Continued approval for metastatic BCC may be contingent upon verification and description of clinical benefit.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Libtayo is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Cutaneous Squamous Cell Carcinoma (must meet all):
 - 1. Diagnosis of CSCC;
 - 2. Disease is metastatic or locally advanced;

CLINICAL POLICY Cemiplimab-rwlc



- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years;
- 5. Member is not a candidate for curative surgery or curative radiation;
- 6. Request meets one of the following (a or b):*
 - a. Dose does not exceed both of the following (i and ii):
 - i. 350 mg every 3 weeks;
 - ii. 1 vial every 3 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
 *Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

B. Basal Cell Carcinoma (must meet all):

- 1. Diagnosis of BCC;
- 2. Disease is metastatic or locally advanced;
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years;
- Previous treatment with a hedgehog pathway inhibitor (e.g., Erivedge[®], Odomzo[®]), unless clinically significant adverse effects are experienced, all are contraindicated, or medical justification indicates that hedgehog pathway inhibitor therapy is not appropriate;
- 6. Request meets one of the following (a or b):*
 - a. Dose does not exceed both of the following (i and ii):
 - i. 350 mg every 3 weeks;
 - ii. 1 vial every 3 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
 - *Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial – 6 months or to the member's renewal date, whichever is longer

C. Non-Small Cell Lung Cancer (must meet all):

- 1. Diagnosis of NSCLC;
- 2. Disease is metastatic or locally advanced where members are not candidates for surgical resection or definitive chemoradiation;
- 3. Prescribed by or in consultation with an oncologist;
- 4. Age \geq 18 years;
- 5. Disease is EGFR wild-type, ALK negative, and ROS1 negative;
- 6. Prescribed in one of the following ways (a, b, or c):
 - a. As a single agent, and one of the following (i or ii):
 - i. Tumor has high PD-L1 expression (TPS \geq 50%);
 - ii. Tumor has PD-L1 expression < 50%, and therapy is prescribed following first-line therapy with Libtayo combination therapy (e.g., cemiplimab-rwlc, [pemetrexed or paclitaxel], and [carboplatin or cisplatin]);

CLINICAL POLICY Cemiplimab-rwlc



- b. In combination with platinum-based chemotherapy (e.g., cisplatin carboplatin);
- c. In combination with pemetrexed as continuation maintenance therapy following first-line therapy with Libtayo combination therapy for nonsquamous cell tumors;
- 7. Request meets one of the following (a or b):*
 - a. Dose does not exceed both of the following (i or ii):
 - i. 350 mg every 3 weeks;
 - ii. 1 vial every 3 weeks;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
 *Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 6 months

Commercial - 6 months or to the member's renewal date, whichever is longer

D. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

- A. All Indications in Section I (must meet all):
 - 1. Currently receiving medication via Centene benefit, or documentation supports that member is currently receiving Libtayo for a covered indication and has received this medication for at least 30 days;
 - 2. Member is responding positively to therapy;
 - 3. If request is for a dose increase, request meets one of the following (a or b):*
 - a. New dose does not exceed both of the following (i or ii):
 - i. 350 mg every 3 weeks;
 - ii. 1 vial every 3 weeks;
 - b. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
 *Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 12 months



Commercial - 6 months or to the member's renewal date, whichever is longer

B. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key ALK: anaplastic lymphoma kinase BCC: basal cell carcinoma CSCC: cutaneous squamous cell carcinoma EGFR: epidermal growth factor receptor

FDA: Food and Drug Administration NSCLC: non-small cell lung cancer PD-1: programmed death receptor-1 TPS: tumor proportion score

Appendix B: Therapeutic Alternatives Not applicable

Appendix C: Contraindications/Boxed Warnings None reported

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
BCC, CSCC, NSCLC	350 mg IV over 30 minutes every 3 weeks	See dosing regimen

VI. Product Availability

Single-dose vial for injection: 350 mg/7 mL (50 mg/mL) solution



VII. References

- 1. Libtayo Prescribing Information. Tarrytown, NY: Regeneron Pharmaceuticals, Inc.; November 2022. Available at: https://www.libtayohcp.com. Accessed November 30, 2022.
- 2. Cemiplimab. In: National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: http://www.nccn.org/professionals/drug_compendium. Accessed December 12, 2022.
- 3. National Comprehensive Cancer Network. Non-Small Cell Lung Cancer, Version 6.2022. Available at https://www.nccn.org/professionals/physician_gls/pdf/nscl.pdf. Accessed December 12, 2022.

Coding Implications

Codes referenced in this clinical policy are for informational purposes only. Inclusion or exclusion of any codes does not guarantee coverage. Providers should reference the most up-todate sources of professional coding guidance prior to the submission of claims for reimbursement of covered services.

HCPCS Codes	Description
J9119	Injection, cemiplimab-rwlc, 1 mg

Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created	10.16.18	11.18
4Q 2019 annual review: no significant changes; references	08.14.19	11.19
reviewed and updated.		
4Q 2020 annual review: modified HIM-Medical Benefit to HIM	10.20.20	11.20
line of business; added HCPCS codes; no significant changes;		
references reviewed and updated.		
RT4: added new indications for BCC and NSCLC; revised	02.24.21	05.21
reference to HIM off-label use policy from HIM.PHAR.21 to		
HIM.PA.154.		
4Q 2021 annual review: no significant changes; references	08.11.21	11.21
reviewed and updated.		
4Q 2022 annual review: no significant changes; references	07.07.22	11.22
reviewed and updated. Template changes applied to other		
diagnoses/indications.		
RT4: added new indication for NSCLC in combination with	12.06.22	
platinum-based chemotherapy; updated criteria per NCCN NSCLC		
guidelines; references reviewed and updated.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional

CLINICAL POLICY Cemiplimab-rwlc



organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.



©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation.