

Clinical Policy: Rimegepant (Nurtec ODT)

Reference Number: CP.PHAR.490

Effective Date: 09.01.20 Last Review Date: 11.22

Line of Business: Commercial, HIM, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Rimegepant (Nurtec[®] [orally disintegrating tablet] ODT) is a calcitonin gene-related peptide receptor (CGRP) antagonist.

FDA Approved Indication(s)

Nurtec ODT is indicated for the:

- Acute treatment of migraine with or without aura in adults
- Preventive treatment of episodic migraine in adults.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Nurtec ODT is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

A. Acute Migraine Treatment (must meet all):

- 1. Diagnosis of migraine headache;
- 2. Age \geq 18 years;
- 3. Failure of at least TWO formulary 5HT_{1B/1D}-agonist migraine medications* (e.g., sumatriptan, rizatriptan, zolmitriptan) at up to maximally indicated doses, unless clinically significant adverse effects are experienced or all are contraindicated; *Prior authorization may be required.
- 4. Failure of Ubrelvy [™]* (at up to maximally indicated doses), unless contraindicated or clinically significant adverse effects are experienced; *Prior authorization may be required.
- 5. For dose increase requests to quantities > 1 box of 8 ODTs per month, member must meet criteria in *Section I, B* below for migraine prophylaxis;
- 6. Nurtec ODT is not prescribed concurrently with other CGRP inhibitors (e.g., Aimovig[®], Ajovy[®], Emgality[®], Qulipta[™], Ubrelvy[®], Vyepti[™]);
- 7. Dose does not exceed 75 mg (1 ODT) per day (one blister pack per month).

Approval duration: 6 months

B. Migraine Prophylaxis (must meet all):

- 1. Diagnosis of episodic migraine;
- 2. Member experiences ≥ 4 migraine days per month for at least 3 months;



- 3. Member does not have chronic migraine, defined as ≥ 15 headache days/month with ≥ 8 migraine days/month for at least 3 months;
- 4. Prescribed by or in consultation with a neurologist, headache, or pain specialist;
- 5. Age \geq 18 years;
- 6. Failure of at least 2 of the following oral migraine preventative therapies, each for 8 weeks and from different therapeutic classes, unless clinically significant adverse effects are experienced or all are contraindicated: antiepileptic drugs (e.g., divalproex sodium, sodium valproate, topiramate), beta-blockers (e.g., metoprolol, propranolol, timolol), antidepressants (e.g., amitriptyline, venlafaxine);
- 7. Failure of at least 1 injectable CGRP therapy (e.g., Aimovig, Ajovy, Emgality, Vyepti), unless clinically significant adverse effects are experienced or all are contraindicated;
- 8. Failure of Qulipta (at up to maximally indicated doses), unless contraindicated or clinically significant adverse effects are experienced;
- 9. If currently receiving treatment with Botox® for migraine prophylaxis and request is for concurrent use of Botox and Nurtec ODT (i.e., not switching from one agent to another), all of the following (a, b, and c):
 - a. Sufficient evidence is provided from at least two high-quality*, published studies in reputable peer-reviewed journals or evidence-based clinical practice guidelines that provide all of the following (i iv):

*Case studies or chart reviews are not considered high-quality evidence

- i. Adequate representation of the member's clinical characteristics, age, and diagnosis;
- ii. Adequate representation of the prescribed drug regimen;
- iii. Clinically meaningful outcomes such as a reduction in monthly migraine or headache days;
- iv. Appropriate experimental design and method to address research questions (see Appendix E for additional information);
- b. Member has experienced and maintained positive response to Botox monotherapy as evidenced by a \geq 30% reduction in migraine days per month from baseline following at least 2 quarterly injection (6 months) of Botox monotherapy;
- c. Despite Botox monotherapy, member continues to experience ≥ 4 migraine days per month and/or severe migraine headaches that result in disability and functional impairment;
- 10. Nurtec ODT is not prescribed concurrently with other CGRP inhibitors (e.g., Aimovig, Ajovy, Emgality, Qulipta, Ubrelvy, Vyepti);
- 11. Dose does not exceed 75 mg (1 ODT) every other day (two blister packs per month). **Approval duration: 3 months**

C. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:



- CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and CP.PMN.255 for Medicaid; or
- b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

A. Acute Migraine Treatment (must meet all):

- 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B);
- 2. Member is responding positively to therapy;
- 3. For dose increase requests to quantities > 1 box of 8 ODTs per month, member must meet criteria in *Section I, B* above for migraine prophylaxis;
- 4. Nurtec ODT is not prescribed concurrently with other CGRP inhibitors (e.g., Aimovig, Ajovy, Emgality, Qulipta, Ubrelvy, Vyepti);*

 *This requirement does not apply to CA if member was previously approved via Centene benefit and is currently stable on therapy with both oral and injectable CGRP inhibitors
- 5. If request is for a dose increase, new dose does not exceed 75 mg (1 ODT) per day (one blister pack per month)

Approval duration: 12 months

B. Migraine Prophylaxis (must meet all):

- 1. Member meets one of the following (a or b):
 - a. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - b. Member is currently receiving medication and is enrolled in a state and product with continuity of care regulations (refer to state specific addendums for CC.PHARM.03A and CC.PHARM.03B);
- 2. Member has experienced and maintained positive response to therapy as evidenced by a reduction in migraine days per month from baseline;
- 3. Nurtec ODT is not prescribed concurrently with other CGRP inhibitors (e.g., Aimovig, Ajovy, Emgality, Qulipta, Ubrelvy, Vyepti);*

 *This requirement does not apply to CA if member was previously approved via Centene benefit and is currently stable on therapy with both oral and injectable CGRP inhibitors
- 4. If request is for a dose increase, new dose does not exceed 75 mg (1 ODT) every other day (two blister packs per month).

Approval duration: 6 months



C. Other diagnoses/indications (must meet 1 or 2):

- 1. If this drug has recently (within the last 6 months) undergone a label change (e.g., newly approved indication, age expansion, new dosing regimen) that is not yet reflected in this policy, refer to one of the following policies (a or b):
 - a. For drugs on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the no coverage criteria policy for the relevant line of business:
 CP.CPA.190 for commercial, HIM.PA.33 for health insurance marketplace, and
 CP.PMN.255 for Medicaid; or
 - b. For drugs NOT on the formulary (commercial, health insurance marketplace) or PDL (Medicaid), the non-formulary policy for the relevant line of business: CP.CPA.190 for commercial, HIM.PA.103 for health insurance marketplace, and CP.PMN.16 for Medicaid; or
- 2. If the requested use (e.g., diagnosis, age, dosing regimen) is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized) AND criterion 1 above does not apply, refer to the off-label use policy for the relevant line of business: CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.

IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key

5-HT: serotonin

AAN: American Academy of Neurology
AHS: American Headache Society

CGRP: calcitonin gene-related peptide
FDA: Food and Drug Administration
ODT: orally disintegrating tablet

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/Maximum Dose			
Abortive Migraine Therapy					
Triptans					
naratriptan (Amerge®)	One tablet (1 or 2.5 mg) PO at	5 mg/day			
	onset; can be repeated in 4 hours				
almotriptan (Axert®)	6.25 to 12.5 mg PO QD	25 mg/day			
	May repeat dose in 2 hours				
frovatriptan (Frova®)	2.5 mg PO QD	7.5 mg/day			
	May repeat dose in 2 hours				



Drug Name	Dosing Regimen	Dose Limit/Maximum
		Dose
sumatriptan (Imitrex® nasal spray)	One spray (5 to 20 mg) at onset into one nostril; can be repeated in 2 hours	40 mg/day
sumatriptan (Imitrex®)	One tablet (25 to 100 mg) PO at onset; can be repeated in two hours	200 mg/day
rizatriptan (Maxalt® /Maxalt MLT®)	One tablet (5 or 10 mg) PO at onset of migraine headache; can be repeated in two hours	30 mg/day
eletriptan (Relpax®)	20 or 40 mg PO QD May repeat dose in 2 hours	40 mg/dose 80 mg/day
zolmitriptan (Zomig [®] /Zomig [®] ZMT)	1.25 or 2.5 mg PO QD May repeat dose in 2 hours	5 mg/dose 10 mg/day
Ubrelvy [™] (ubrogepant)	50 or 100 mg PO, as needed. If needed, a second dose may be administered at least 2 hours after the initial dose. The maximum dose in a 24-hour period is 200 mg.	200 mg/day
	Prophylactic Migraine Therapy	
Anticonvulsants such as: divalproex (Depakote®), topiramate (Topamax®), valproate sodium	Migraine Prophylaxis Refer to prescribing information or Micromedex	Refer to prescribing information or Micromedex
Beta-blockers such as: propranolol (Inderal®), metoprolol (Lopressor®)*, timolol, atenolol (Tenormin®)*, nadolol (Corgard®)*	Migraine Prophylaxis Refer to prescribing information or Micromedex	Refer to prescribing information or Micromedex
Antidepressants/tricycli c antidepressants* such as: amitriptyline (Elavil®), venlafaxine (Effexor®)	Migraine Prophylaxis Refer to prescribing information or Micromedex	Refer to prescribing information or Micromedex
Qulipta [™] (atogepant)	10 mg, 30 mg, or 60 mg PO QD	60 mg/day



Drug Name	Dosing Regimen	Dose Limit/Maximum Dose
Aimovig [™] (erenumabaooe)	70 mg SC once monthly	140 mg/month
,	Some patients may benefit from a dosage of 140 mg injected subcutaneously once monthly	
Ajovy® (fremanezumab-vfrm)	225 mg SC once monthly or 675 mg SC every three months	675 mg every 3 months
Emgality® (galcanezumab-gnlm)	Loading dose: 240 mg SC once Maintenance dose: 120 mg SC once monthly	120 mg/month
Vyepti [™] (eptinezumab- jjmr)	The recommended dosage is 100 mg IV every 3 months.	300 mg every 3 months
	Some patients may benefit from a dosage of 300 mg IV every 3 months.	

Therapeutic alternatives are listed as Brand name® (generic) when the drug is available by brand name only and generic (Brand name®) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): history of hypersensitivity reaction to rimegepant, Nurtec ODT, or to any of its components.
- Boxed warning(s): none reported

Appendix D: General Information

The American Headache Society (2018) provides the following migraine guidance:

- Migraine patients who need to use acute treatments on a regular basis should be instructed to limit treatment to an average of 2 headache days per week, and patients observed to be exceeding this limit should be offered preventive treatment. Indications for preventive treatment:
 - o Attacks significantly interfere with patients' daily routines despite acute treatment
 - o Frequent attacks (≥ 4 migraine headache days [per month])
 - o Contraindication to, failure, or overuse of acute treatments, with overuse defined as:
 - 10 or more days per month for ergot derivatives, triptans, opioids, combination analgesics, and a combination of drugs from different classes that are not individually overused
 - 15 or more days per month for non-opioid analgesics, acetaminophen, and nonsteroidal anti-inflammatory drugs (NSAIDs [including aspirin])
 - Adverse effects with acute treatments
 - Patient preference



 Prevention should also be considered in the management of certain uncommon migraine subtypes, including hemiplegic migraine, migraine with brainstem aura, migraine with prolonged aura, and those who have previously experienced a migrainous infarction, even if there is low attack frequency.

Appendix E: Appropriate Experimental Design Methods

- Randomized, prospective controlled trials are generally considered the gold standard; however:
 - o In some clinical studies, it may be unnecessary or not feasible to use randomization, double-blind trials, placebos, or crossover.
 - Non-randomized prospective clinical trials with a significant number of subjects may be a basis for supportive clinical evidence for determining accepted uses of drugs.
- Case reports and chart reviews are generally considered uncontrolled and anecdotal information and do not provide adequate supportive clinical evidence for determining accepted uses of drugs.

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Migraine -	75 mg PO as needed. The maximum dose in a 24-hour	75 mg/day
acute	period is 75 mg. The safety of using more than 18 doses	
treatment	in a 30-day period has not been established.	
Migraine	75 mg PO every other day	75 mg/dose
prophylaxis	-	_

VI. Product Availability

ODT (blister pack of 8): 75 mg

VII. References

- 1. Nurtec ODT Prescribing Information. New Haven, CT: Biohaven Pharmaceuticals, Inc.; April 2022. Available at https://biohaven-nurtec-consumer-assets.s3.amazonaws.com/nurtec-prescribing-information.pdf. Accessed July 27, 2022.
- 2. Croop R, Goadsby PJ, Stock DA, et al. Efficacy, safety, and tolerability of rimegepant orally disintegrating tablet for the acute treatment of migraine: a randomised, phase 3, double-blind, placebo-controlled trial. The Lancet. August 31, 2019; 394:737-745.
- 3. MICROMEDEX® Healthcare Series [Internet database]. Greenwood Village, Colo: Thomson Healthcare. Updated periodically. Accessed July 27, 2022.
- 4. American Headache Society. The American Headache Society position statement on integrating new migraine treatments into clinical practice. Headache. 2019;59:1-18.
- 5. Silberstein SD, Holland S, Freitag F, Dodick DW, Argoff C, Ashman E. Evidence-based guideline update: Pharmacologic treatment for episodic migraine prevention in adults: Report of the Quality Standards Subcommittee of the American Academy of Neurology and the American Headache Society. Neurology. 2012;78:1337-1345.
- 6. Croop R, Lipton RB, Kudrow D, et al. Oral rimegepant for preventive treatment of migraine: a phase 2/3, randomised, double-blind, placebo-controlled trial. Lancet 2021; 397: 51–60.



Reviews, Revisions, and Approvals	Date	P&T Approval
Deliver courts 1	04.14.20	Date
Policy created	04.14.20	08.20
3Q 2021 annual review: RT4 added new indication for episodic	06.28.21	08.21
migraine prophylaxis; added clarification in continuation of therapy		
to indicate requirement for concurrent use with other CGRP		
inhibitors does not apply to CA if member was previously		
approved via Centene benefit and is currently stable on therapy		
with both oral and injectable CGRP inhibitors; replaced		
HIM.PHAR.21 to reference HIM.PA.154; references reviewed and		
updated.		
1Q 2022 annual review: per SDC and prior clinical guidance for	11.17.21	02.22
migraine prophylaxis added redirection to newly approved oral		
CGRP Qulipta; references reviewed and updated.		
4Q 2022 annual review: Added criteria for concurrent use with	08.23.22	11.22
Botox requiring supportive evidence from published studies or		
clinical practice guidelines, positive response with Botox		
monotherapy, and continued migraine burden; per August SDC and		
prior clinical guidance for migraine prophylaxis added redirection		
to injectable CGRP, for acute migraine treatment added redirection		
to Ubrelvy; references reviewed and updated. Template changes		
applied to other diagnoses/indications and continued therapy		
section.		

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.



This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2020 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene® and Centene Corporation® are registered trademarks exclusively owned by Centene Corporation.