

Clinical Policy: Hydroxyurea (Siklos)

Reference Number: CP.PMN.193 Effective Date: 02.19.19 Last Review Date: 05.21 Line of Business: Commercial, HIM, Medicaid

Revision Log

See <u>Important Reminder</u> at the end of this policy for important regulatory and legal information.

Description

Hydroxyurea (Siklos[®]) is an antimetabolite.

FDA Approved Indication(s)

Siklos is indicated to reduce the frequency of painful crises and to reduce the need for blood transfusions in pediatric patients, 2 years of age and older, with sickle cell anemia with recurrent moderate to severe painful crises.

Policy/Criteria

Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.

It is the policy of health plans affiliated with Centene Corporation[®] that Siklos is **medically necessary** when the following criteria are met:

I. Initial Approval Criteria

- A. Sickle Cell Disease (must meet all):
 - 1. Diagnosis of sickle cell disease;
 - 2. Age \geq 2 years;
 - 3. Member must use generic hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced;
 - 4. Dose does not exceed 35 mg per kg per day based on weight.

Approval duration:

Medicaid/HIM – 12 months

Commercial – Length of Benefit

B. Oncology Indications (off-label) (must meet all):

- 1. Diagnosis of one of the following (a, b, c, or d);
 - a. Acute myeloid leukemia;
 - b. Chronic myeloid leukemia;
 - c. Head and neck cancer;
 - d. Myeloproliferative neoplasms (myelofibrosis, polycythemia vera, essential thrombocythemia);
 - e. Myelodysplastic syndromes;
- 2. Age \geq 2 years;
- 3. Member must use generic hydroxyurea, unless contraindicated or clinically significant adverse effects are experienced;

CLINICAL POLICY Hydroxyurea



- 4. Request meets one of the following (a or b):
 - a. Dose does not exceed 80 mg per kg per day based on weight;
 - b. Dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).

Approval duration:

Medicaid/HIM – 12 months **Commercial** – Length of Benefit

C. Other diagnoses/indications

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

II. Continued Therapy

- A. All Indications in Section I (must meet all):
 - 1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;
 - 2. Member is responding positively to therapy;
 - 3. If request is for a dose increase, request meets one of the following (a, b, or c):
 - a. Sickle cell disease: new dose does not exceed 35 mg per kg per day based on weight;
 - b. Oncology indications: new dose does not exceed 80 mg per kg per day based on weight;*
 - c. New dose is supported by practice guidelines or peer-reviewed literature for the relevant off-label use (*prescriber must submit supporting evidence*).
 *Prescribed regimen must be EDA gunnaued on recommended by NCCN

*Prescribed regimen must be FDA-approved or recommended by NCCN

Approval duration:

Medicaid/HIM – 12 months **Commercial** – Length of Benefit

B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

Approval duration: Duration of request or 6 months (whichever is less); or

 Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid.

III. Diagnoses/Indications for which coverage is NOT authorized:

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – CP.CPA.09 for commercial, HIM.PA.154 for health insurance marketplace, and CP.PMN.53 for Medicaid, or evidence of coverage documents.



IV. Appendices/General Information

Appendix A: Abbreviation/Acronym Key FDA: Food and Drug Administration

Appendix B: Therapeutic Alternatives

This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
hydroxyurea	Sickle cell disease: 15 mg/kg PO QD	Sickle disease: 35
(Hydrea [®] ,	CML: 40 mg/kg/day	mg/kg/day
Droxia [®])	Head and neck cancer: 1,000 mg q12h	Oncology indications: 80
,		mg/kg/dav

Therapeutic alternatives are listed as Brand name[®] (generic) when the drug is available by brand name only and generic (Brand name[®]) when the drug is available by both brand and generic.

Appendix C: Contraindications/Boxed Warnings

- Contraindication(s): hypersensitivity to hydroxyurea or any other component of its formulation
- Boxed warning(s): myelosuppression and malignancies

V. Dosage and Administration

Indication	Dosing Regimen	Maximum Dose
Sickle cell	Initial dose 20 mg/kg PO QD. Dose may	35 mg/kg/day (maximum
disease	be increased by 5 mg/kg/day every 8	dose based on weight)
	weeks or sooner if a severe painful crisis	
	occurs.	

VI. Product Availability

Tablets: 100 mg, 1,000 mg

VII. References

- 1. Siklos Prescribing Information. Bryn Mawr, PA: Medunik USA, Inc.; May 2019. Available at <u>https://files.medunik.com/usa/siklos/prescribing-information.pdf</u>. Accessed February 1, 2021.
- 2. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2016. Available at: <u>http://www.clinicalpharmacology-ip.com</u>. Accessed February 13, 2020.
- 3. Lexicomp Online [Internet Database]. Hudson, Ohio: Wolters Kluwer Clinical Drug Information, Inc. Updated periodically. Accessed February 13, 2020.
- 4. National Comprehensive Cancer Network Drugs and Biologics Compendium. Available at: <u>http://www.nccn.org/professionals/drug_compendium</u>. Accessed February 12, 2020.
- Brandow A, Carroll C, Creary S, et al. American Society of Hematology 2020 guidelines for sickle cell disease: management of acute and chronic pain. *Blood Advances*. 2020;4(12):2656-2701.



Reviews, Revisions, and Approvals	Date	P&T Approval Date
Policy created per SDC.	02.19.19	05.19
2Q 2020 annual review: no significant changes; applied HIM line of business; references reviewed and updated	02.13.20	05.20
2Q 2021 annual review: myelodysplastic syndromes added as option for off-label oncology indication per NCCN-supported category 2A recommendation; references for HIM line of business off-label use revised from HIM.PHAR.21 to HIM.PA.154; references reviewed and updated.	02.01.21	05.21

Important Reminder

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. "Health Plan" means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan's affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to

CLINICAL POLICY Hydroxyurea



recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

Note:

For Medicaid members, when state Medicaid coverage provisions conflict with the coverage provisions in this clinical policy, state Medicaid coverage provisions take precedence. Please refer to the state Medicaid manual for any coverage provisions pertaining to this clinical policy.

©2019 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene[®] and Centene Corporation[®] are registered trademarks exclusively owned by Centene Corporation.