

## **Clinical Policy: Insulin Glargine (Rezvoglar, Semglee, Toujeo)**

Reference Number: HIM.PA.09

Effective Date: 03.01.19

Last Review Date: 02.22

Line of Business: HIM

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

### **Description**

Insulin glargine (Rezvoglar<sup>™</sup>, Semglee<sup>®</sup>, Toujeo<sup>®</sup>) is a long-acting human insulin analogs.

### **FDA Approved Indication(s)**

Rezvoglar and Semglee are indicated to improve glycemic control in adults and pediatric patients with type 1 diabetes mellitus and in adults with type 2 diabetes mellitus.

Toujeo is indicated to improve glycemic control in adults and pediatric patients 6 years and older with diabetes mellitus.

Limitation(s) of use: Rezvoglar, Semglee and Toujeo are not recommended for treating diabetic ketoacidosis.

### **Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Rezvoglar, Semglee and Toujeo are **medically necessary** when the following criteria are met:

#### **I. Initial Approval Criteria**

##### **A. Diabetes Mellitus (must meet all):**

1. Diagnosis of type 1 or type 2 diabetes mellitus;
2. Age  $\geq$  6 years;
3. Failure of Basaglar<sup>®</sup>, Levemir<sup>®</sup>, and Tresiba<sup>®</sup> unless clinically significant adverse effects are experienced or all are contraindicated.

**Approval duration: 12 months**

##### **B. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PA.154 for health insurance marketplace.

#### **II. Continued Therapy**

##### **A. Diabetes Mellitus (must meet all):**

1. Currently receiving medication via Centene benefit or member has previously met initial approval criteria;

2. Member is responding positively to therapy.

**Approval duration: 12 months**

**B. Other diagnoses/indications (must meet 1 or 2):**

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

**Approval duration: Duration of request or 12 months (whichever is less); or**

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PA.154 for health insurance marketplace.

**III. Diagnoses/Indications for which coverage is NOT authorized:**

A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policies – HIM.PA.154 for health insurance marketplace or evidence of coverage documents.

**IV. Appendices/General Information**

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

<b>Drug Name</b>	<b>Dosing Regimen</b>	<b>Dose Limit/ Maximum Dose</b>
Basaglar <sup>®</sup> (insulin glargine)	<p><b>Type 1 diabetes mellitus:</b> Initiation: Approximately one-third of the total daily insulin requirement administered SC QD</p> <p><b>Type 2 diabetes mellitus:</b> Initiation: 0.2 units/kg SC QD up to 10 units/day. Adjust dosage according to patient response</p>	Not applicable
Levemir <sup>®</sup> (insulin detemir)	Individualize starting dose based on type of diabetes and whether patient is insulin-naïve. Administer SC QD (with evening meal or at bedtime) or BID.	Not applicable
Tresiba <sup>®</sup> (insulin degludec)	<p><b>Type 1 diabetes mellitus:</b> Initiation:</p> <ul style="list-style-type: none"> <li>• Insulin-naïve: 1/3 to 1/2 of total daily insulin dose SC QD</li> <li>• Already on insulin: SC QD: <ul style="list-style-type: none"> <li>○ Adults: same unit dose as total daily long or intermediate-acting insulin unit dose</li> <li>○ Pediatrics: 80% of total daily long or intermediate-acting insulin unit dose</li> </ul> </li> </ul>	Not applicable

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	<b>Type 2 diabetes mellitus:</b> Initiation: <ul style="list-style-type: none"> <li>• Insulin-naïve: 10 units SC QD</li> <li>• Already on insulin: SC QD:               <ul style="list-style-type: none"> <li>○ Adults: same unit dose as total daily long or intermediate-acting insulin unit dose</li> <li>○ Pediatrics: 80% of total daily long or intermediate-acting insulin unit dose</li> </ul> </li> </ul>	

Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.

*Appendix C: Contraindications/Boxed Warnings*

- Contraindication(s): use during episodes of hypoglycemia, hypersensitivity to the requested product or one of its excipients
- Boxed warning(s): none reported

**V. Dosage and Administration**

Drug Name	Indication	Dosing Regimen	Maximum Dose
Insulin glargine (Toujeo)	Type 1 diabetes mellitus	Initiation: <ul style="list-style-type: none"> <li>• Insulin-naïve: 1/3 to 1/2 of total daily insulin dose SC QD</li> <li>• Already on insulin: SC QD:               <ul style="list-style-type: none"> <li>○ Once-daily long or intermediate insulin: same unit dose as total daily long acting insulin unit dose. Expect higher daily dose of Toujeo will be needed to maintain the same level of glycemic control in patients on Lantus.</li> <li>○ Twice-daily long or intermediate insulin: 80% of total daily long or intermediate-acting insulin unit dose</li> </ul> </li> </ul>	Not applicable; dose is individualized and titrated based on metabolic needs, blood glucose monitoring results, and glycemic control goal
	Type 2 diabetes mellitus	Initiation: <ul style="list-style-type: none"> <li>• Insulin-naïve: 0.2 units/kg SC QD</li> <li>• Already on insulin: SC QD:               <ul style="list-style-type: none"> <li>○ Once-daily long or intermediate insulin: same unit dose as total daily long acting insulin unit dose. Expect higher daily dose of Toujeo will be needed to maintain the same level of</li> </ul> </li> </ul>	Not applicable; dose is individualized and titrated based on metabolic needs, blood glucose monitoring results, and glycemic control goal

Drug Name	Indication	Dosing Regimen	Maximum Dose
		glycemic control in patients on Lantus. <ul style="list-style-type: none"> <li>Twice-daily long or intermediate insulin: 80% of total daily long or intermediate-acting insulin unit dose</li> </ul>	
Insulin glargine-yfgn (Semglee), Insulin glargine-aglr (Rezvoglar)	Type 1 diabetes mellitus	Initiation: Approximately one-third of the total daily insulin requirement administered SC QD	Not applicable
	Type 2 diabetes mellitus	Initiation: 0.2 units/kg SC QD or 10 units/day. Adjust dosage according to patient response	Not applicable

#### VI. Product Availability

Drug Name	Availability
Insulin glargine (Toujeo)	Single-patient-use prefilled pen 300 units/mL: 1.5 mL (Toujeo SoloStar), 3 mL (Toujeo Max SoloStar)
Insulin glargine-yfgn (Semglee)	<ul style="list-style-type: none"> <li>Multiple-dose vial: 10 mL containing 100 units/mL</li> <li>Prefilled pen: 3 mL containing 100 units/mL</li> </ul>
Insulin glargine-aglr (Rezvoglar)	KwikPen <sup>®</sup> prefilled pen: 3 mL containing 100 units/mL

#### VII. References

- Semglee Prescribing Information. Morgantown, WV: Mylan Specialty L.P.; June 2020. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2020/210605s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2020/210605s000lbl.pdf). Accessed September 10, 2021.
- Toujeo Prescribing Information. Bridgewater, NJ: Sanofi\_Aventis U.S. LLC; December 2020. Available at: [www.toujeo.com](http://www.toujeo.com). Accessed October 28, 2021.
- Rezvoglar Prescribing Information. Indianapolis, IN: Eli Lilly and Company; December 2021. Available at: [https://www.accessdata.fda.gov/drugsatfda\\_docs/label/2021/761215s000lbl.pdf](https://www.accessdata.fda.gov/drugsatfda_docs/label/2021/761215s000lbl.pdf). Accessed January 10, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
1Q 2019 Policy created	12.05.18	02.19
Per SDC and prior clinical guidance, added additional requirement for redirection to Levemir.	10.23.19	

<b>Reviews, Revisions, and Approvals</b>	<b>Date</b>	<b>P&amp;T Approval Date</b>
1Q 2020 annual review: added requirement for trial of Levemir per SDC; references reviewed and updated.	10.24.19	02.20
Added Semglee to policy per October SDC and prior clinical guidance	10.08.20	
1Q 2021 annual review: no significant changes; references to HIM.PHAR.21 revised to HIM.PA.154; references reviewed and updated.	10.26.20	02.21
Per October SDC and prior clinical guidance, removed Tresiba from policy as PA is no longer required; added Toujeo to policy and revised redirection to require use of Basaglar, Levemir, and Tresiba; revised required age to 6 years or older consistent with Semglee and Toujeo prescribing information.	10.27.21	
1Q 2022 annual review: RT4: added Rezvoglar to policy; references reviewed and updated.	01.10.22	02.22

**Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2018 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.