

**Clinical Policy: Spinosad (Natroba)**

Reference Number: HIM.PA.134

Effective Date: 12.01.17

Last Review Date: 08.22

Line of Business: HIM

[Revision Log](#)

See [Important Reminder](#) at the end of this policy for important regulatory and legal information.

**Description**

Spinosad topical suspension (Natroba<sup>®</sup>) is a pediculicide and scabicide.

**FDA Approved Indication(s)**

Natroba is indicated for the topical treatment of:

- Head lice infestations in patients 6 months of age and older
- Scabies infestations in adult and pediatric patients 4 years of age and older.

**Policy/Criteria**

*Provider must submit documentation (such as office chart notes, lab results or other clinical information) supporting that member has met all approval criteria.*

It is the policy of health plans affiliated with Centene Corporation<sup>®</sup> that Natroba is **medically necessary** when the following criteria are met:

**I. Initial Approval Criteria****A. Head Lice** (must meet all):

1. Diagnosis of head lice;
2. Age  $\geq$  6 months;
3. Failure of permethrin 1% cream in the last 60 days, unless contraindicated or clinically significant adverse effects are experienced;
4. Request does not exceed 2 bottles (8 oz).

**Approval duration: 14 days**

**B. Scabies Infestation** (must meet all):

1. Diagnosis of scabies infestation;
2. Age  $\geq$  4 years;
3. Failure of permethrin 5% cream, unless contraindicated or clinically significant adverse effects are experienced;
4. Request does not exceed 4 bottles (16 oz).

**Approval duration: one time**

**C. Other diagnoses/indications**

1. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PA.154 for health insurance marketplace.

## II. Continued Therapy

### A. All Indications in Section I (must meet all):

1. Re-authorization is not permitted. Members must meet the initial approval criteria.

**Approval duration: Not applicable**

### B. Other diagnoses/indications (must meet 1 or 2):

1. Currently receiving medication via Centene benefit and documentation supports positive response to therapy.

**Approval duration: Duration of request or 14 days (whichever is less); or**

2. Refer to the off-label use policy for the relevant line of business if diagnosis is NOT specifically listed under section III (Diagnoses/Indications for which coverage is NOT authorized): HIM.PA.154 for health insurance marketplace.

## III. Diagnoses/Indications for which coverage is NOT authorized:

- A. Non-FDA approved indications, which are not addressed in this policy, unless there is sufficient documentation of efficacy and safety according to the off label use policy – HIM.PA.154 for health insurance marketplace or evidence of coverage documents.

## IV. Appendices/General Information

*Appendix A: Abbreviation/Acronym Key*

FDA: Food and Drug Administration

*Appendix B: Therapeutic Alternatives*

*This table provides a listing of preferred alternative therapy recommended in the approval criteria. The drugs listed here may not be a formulary agent for all relevant lines of business and may require prior authorization.*

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
permethrin 1% creme rinse/lotion	Adults, adolescents, children, and infants $\geq 2$ months: Shampoo hair with regular shampoo, rinse and towel dry. Then, apply permethrin 1% lotion sufficient to saturate the hair and scalp (usually 25 to 30 mL), especially behind the ears and on the nape of the neck. Leave on hair for 10 minutes but no longer. Then, rinse thoroughly with water. If live lice are seen 7 days or more after the first application, a second treatment should be given.	One application to affected area
permethrin 5% cream	Thoroughly massage permethrin 5% cream into the skin from the head to the soles of the feet. Scabies rarely infests the scalp of adults, although the hairline, neck, temple, and forehead may be infested in infants and geriatric patients. Usually 30 grams is sufficient for an average adult. The cream should be removed by washing (shower or bath) after 8 to 14 hours. Infants should be treated	One application to affected area

Drug Name	Dosing Regimen	Dose Limit/ Maximum Dose
	on the scalp, temple, and forehead. One application is generally curative. Patients may experience persistent pruritus after treatment. This is rarely a sign of treatment failure and is not an indication for retreatment. Demonstrable living mites after 14 days indicate that retreatment is necessary.	

*Therapeutic alternatives are listed as Brand name<sup>®</sup> (generic) when the drug is available by brand name only and generic (Brand name<sup>®</sup>) when the drug is available by both brand and generic.*

*Appendix C: Contraindications/Boxed Warnings*

None reported

**V. Dosage and Administration**

Indication	Dosing Regimen	Maximum Dose
Head lice	Apply a sufficient amount to cover dry scalp, then apply to dry hair. Depending on hair length, apply up to 120 mL (one bottle) to adequately cover scalp and hair. Leave on for 10 minutes, then thoroughly rinse off with warm water. If live lice are seen 7 days after the first treatment, a second treatment should be applied.	120 mL/application
Scabies infestation	Apply a sufficient amount of Natroba to skin to completely cover the body from the neck to the toes (including the soles of the feet). For patients with balding scalp, also apply product to the scalp, hairline, temples, and forehead. Allow to absorb into the skin and dry for 10 minutes before getting dressed. Leave on the skin for at least 6 hours before showering or bathing.	Varies per body surface area

**VI. Product Availability**

Suspension: 9 mg of spinosad per gram of Natroba topical suspension in 120 mL bottles

**VII. References**

1. Natroba Prescribing Information. Carmel, IN: ParaPRO LLC; April 2021. Available at: <http://www.natroba.com>. Accessed March 22, 2022.
2. Centers for Disease Control and Prevention. Parasites-Lice-Head Lice. Available at: <https://www.cdc.gov/parasites/lice/head/treatment.html>. Updated October 15, 2019. Accessed March 22, 2022.
3. Centers for Disease Control and Prevention. Parasites - Scabies. Suggested General Guidelines. Available at: <http://www.cdc.gov/parasites/scabies/treatment.html>. September 3, 2015; Accessed March 22, 2022.

4. Clinical Pharmacology [database online]. Tampa, FL: Gold Standard, Inc.; 2020. Available at: <http://www.clinicalpharmacology-ip.com/>. Accessed March 22, 2022.

Reviews, Revisions, and Approvals	Date	P&T Approval Date
4Q 2018 annual review: modified timeframe of trial of permethrin from within the last 6 months to last 60 days; shortened approval duration from 1 month to 14 days (aligns with other pediculicide policies); continued therapy: removed requirement that 6 months should have elapsed since previous claim for Natroba; references reviewed and updated.	08.08.18	11.18
3Q 2019 annual review: no significant changes; references reviewed and updated.	05.31.19	08.19
3Q 2020 annual review: no significant changes; references reviewed and updated.	08.08.20	08.20
3Q 2021 annual review: added criteria for newly approved indication for scabies infestation; references to HIM.PHAR.21 revised to HIM.PA.154; references reviewed and updated.	05.08.21	08.21
3Q 2022 annual review: no significant changes; references reviewed and updated.	03.22.22	08.22

### **Important Reminder**

This clinical policy has been developed by appropriately experienced and licensed health care professionals based on a review and consideration of currently available generally accepted standards of medical practice; peer-reviewed medical literature; government agency/program approval status; evidence-based guidelines and positions of leading national health professional organizations; views of physicians practicing in relevant clinical areas affected by this clinical policy; and other available clinical information. The Health Plan makes no representations and accepts no liability with respect to the content of any external information used or relied upon in developing this clinical policy. This clinical policy is consistent with standards of medical practice current at the time that this clinical policy was approved. “Health Plan” means a health plan that has adopted this clinical policy and that is operated or administered, in whole or in part, by Centene Management Company, LLC, or any of such health plan’s affiliates, as applicable.

The purpose of this clinical policy is to provide a guide to medical necessity, which is a component of the guidelines used to assist in making coverage decisions and administering benefits. It does not constitute a contract or guarantee regarding payment or results. Coverage decisions and the administration of benefits are subject to all terms, conditions, exclusions and limitations of the coverage documents (e.g., evidence of coverage, certificate of coverage, policy, contract of insurance, etc.), as well as to state and federal requirements and applicable Health Plan-level administrative policies and procedures.

This clinical policy is effective as of the date determined by the Health Plan. The date of posting may not be the effective date of this clinical policy. This clinical policy may be subject to applicable legal and regulatory requirements relating to provider notification. If there is a

discrepancy between the effective date of this clinical policy and any applicable legal or regulatory requirement, the requirements of law and regulation shall govern. The Health Plan retains the right to change, amend or withdraw this clinical policy, and additional clinical policies may be developed and adopted as needed, at any time.

This clinical policy does not constitute medical advice, medical treatment or medical care. It is not intended to dictate to providers how to practice medicine. Providers are expected to exercise professional medical judgment in providing the most appropriate care, and are solely responsible for the medical advice and treatment of members. This clinical policy is not intended to recommend treatment for members. Members should consult with their treating physician in connection with diagnosis and treatment decisions.

Providers referred to in this clinical policy are independent contractors who exercise independent judgment and over whom the Health Plan has no control or right of control. Providers are not agents or employees of the Health Plan.

This clinical policy is the property of the Health Plan. Unauthorized copying, use, and distribution of this clinical policy or any information contained herein are strictly prohibited. Providers, members and their representatives are bound to the terms and conditions expressed herein through the terms of their contracts. Where no such contract exists, providers, members and their representatives agree to be bound by such terms and conditions by providing services to members and/or submitting claims for payment for such services.

©2017 Centene Corporation. All rights reserved. All materials are exclusively owned by Centene Corporation and are protected by United States copyright law and international copyright law. No part of this publication may be reproduced, copied, modified, distributed, displayed, stored in a retrieval system, transmitted in any form or by any means, or otherwise published without the prior written permission of Centene Corporation. You may not alter or remove any trademark, copyright or other notice contained herein. Centene<sup>®</sup> and Centene Corporation<sup>®</sup> are registered trademarks exclusively owned by Centene Corporation.